

AVIAN INFLUENZA VIRUS AND NEWCASTLE VIRUS SURVEILLANCE AND CHARACTERIZATION IN BROILER AND LAYER CHICKEN FLOCKS IN EGYPT

AZHAR G. SHALABY^{*}; AHMED M. ERFAN^{*}; FATMA A. ABDEL REHEEM^{*}; ABDULLAH A. SELIM^{**}; MOHAMED H. AL HUSSENY^{****}; SHEREEN G. KHOLOS^{***} and SOAD A. NASEF^{**}

^{*}Biotechnology Department, Reference Laboratory for Veterinary Quality Control on Poultry Production, Animal Health Research Institute, Nadi El-Seid St., P.O. Box 246, Dokki, Giza 12618, Egypt.

^{**}Poultry Diseases Department, Reference Laboratory for Veterinary Quality Control on Poultry Production, Animal Health Research Institute, Nadi El-Seid St., P.O. Box 246, Dokki, Giza 12618, Egypt.

^{***}Epidemiology Department, Reference Laboratory for Veterinary Quality Control on Poultry Production, Animal Health Research Institute, Nadi El-Seid St., P.O. Box 246, Dokki, Giza 12618, Egypt.

^{****}Biochemistry Department, Reference Laboratory for Veterinary Quality Control on Poultry Production, Animal Health Research Institute, Nadi El-Seid St., P.O. Box 246, Dokki, Giza 12618, Egypt.

Email: azhar_gaber@yahoo.com

ABSTRACT

Received at: 16/6/2014

Accepted: 6/8/2014

Considering that highly pathogenic avian influenza H5N1 endemically circulates in the Egyptian poultry population, Active surveillance was undertaken in 195 broiler and layers farms from Eighteen Egyptian governorates resulted in, 34,7 and 24 positive farms for AI H9N2, AI H5N1 and Virulent ND respectively, using RT-PCR. The genetic analysis of avian influenza viruses of HA gene of the H5N1 type revealed that these viruses are highly pathogenic strains as they are tracking clade 2.2.1 of classic viruses which became endemic in Egypt since 2006. However, H9N2 viruses revealed that they are low pathogenic as they are tracking G1-lineage of Asian viruses which transmitted to Middle East since 2000. Also sequence analysis for F gene for twelve NDV isolates revealed that they are of virulent type and genetically related to genotype VII NDV except two isolates are avirulent type and resemble Lasota-like strains. Pathogenicity studies for NDV isolates using ICPI was evaluated for the strain (NDV-EG-567f-2012) which of virulent with ICPI value 1.96. The analysis of isolated strains by different diagnostic and analytical techniques implies that the H5N1 and H9N2 silent infection exist in Egypt.

Key words: Pathogenicity- Lasota like strains- lineage- RT-PCR- endemic.

INTRODUCTION

The highly pathogenic avian influenza virus subtype H5N1 was first recognized in poultry in Egypt in 2006 (Aly *et al.*, 2008). The subtype H9N2 is widely circulated in the Middle East region and it was associated with serious disease in poultry (Aamir *et al.*, 2007; Alexander, 2007; Perk *et al.*, 2009). In addition, recent detection of low pathogenic avian influenza H9N2 subtype has been recorded in bobwhite quail in Egypt in 2011 with genetic relatedness to the G1-lineage that represented by A/Quail/Hong Kong/G1/97 (Qa/HK/G1/97) that still circulated in the Middle East region. (El-Zoghby *et al.*, 2012) Standard RT-PCR has been applied previously to the detection of AIV and each of the 15 HA subtypes (Starick *et al.*, 2000; Lee *et al.*, 2001; Munch *et al.*, 2001). Sequence analysis of the viral genes and their respective amino acids is a valuable tool for detection of any genic variation among viruses circulating in the field and also provides

information about the source of the initial disease outbreak in a certain geographic district (Duvvuri *et al.*, 2009). The broad variation in virulence and clinical signs necessitates the careful definition of what constitutes ND for the purposes of trade, control measures and policies. The definition of ND currently in use in all member states of the European Union is defined in Directive 92/66/EEC of the Commission for European Communities (OIE 2012). ND outbreaks occur frequently in Egypt and the source of the virulent NDV in these outbreaks are not known. (Mohamed *et al.*, 2009) reported the complete genome sequence of a NDV strain isolated from an outbreak at a poultry farm in Al-Sharkia province, Egypt in 2005 (chicken/Egypt/1/2005).

In this work, the surveillance were carried out to detect the avian influenza and Newcastle viruses and evaluate their virulence attitude as considered most important respiratory viruses for poultry and specially broiler and layer sectors and its zoonotic

picture. Furthermore discuss the genetic characterizations for Newcastle and AI with its subtypes H5N1 and H9N2.

MATERIALS and METHODS

195 layer and broiler farms' from eighteen Egyptian governorates have been survived during late of 2011-2012. 2448 cloacal swabs and 306 organs like trachea and lung were collected and examined as shown in Table. (1).

Table 1: Collective Numbers of farms, samples in relation to governorate and type of production.

No.	Governorates	No. of farms	No of examined samples		Type of production	
			Cloacal swabs	Organs (Tr& lung)	Layers	Broilers
1	Al Fayoum	51	920	65	5	46
2	Al Qalubia	20	150	14	1	19
3	Al Dakahlia	19	294	34	3	16
4	Al Ismalia	18	85	22	0	18
5	Benisueif	15	204	26	3	12
6	Domiatte	11	95	47	2	9
7	Luxor	12	22	9	0	12
8	Al Giza	9	186	10	4	5
9	Al Sharkia	10	180	8	3	7
10	Al Alexandria	8	90	26	4	4
11	Kafr el sheikh	5	46	10	0	5
12	Al Behira	5	64	11	0	5
13	Al Monefia	3	44	2	0	3
14	Al Gharbia	3	20	11	0	3
15	Al Minia	2	38	2	0	2
16	North sinai	2	4	6	0	2
17	Aswan	1	-	1	0	1
18	Port said	1	6	1	0	1
Total		195	2448	306	25	170

Real time PCR:

This test was done for the different samples to detect the AI common gene and haemagglutinin genes of H5 and H9 by multiplex PCR also the method was done to detect the F gene for Newcastle and determine the virulence of the positive strain.

RNA was extracted from samples was performed using the QIAamp viral RNA Mini kit (Qiagen, Germany, GmbH) according to the manufacturer's recommendations. Briefly 140 µl of the sample suspension was incubated with 5.6 µl of carrier RNA and 560 µl of AVL buffer at room temperature for 10 min. After incubation, 560 µl of 100% ethanol was

added to the lysate. The mixture was then transferred to the silica column then centrifuged and washed following the manufacturer's recommendations. Nucleic acid was eluted with 60 µl of elution buffer (AE) provided within the kit] NDV, and AI: Primers were utilized in a 25- µl reaction containing 12.5 µl of Quantitect probe rt-PCR master mix (Qiagen, Germany, GmbH), 0.5 µl of each primer of 50 pmol concentration, 0.125 µl of the specific probe, 4.5 µl of water, 0.25 µl of the rt-enzyme and 6 µl of RNA extract.. The reaction was performed in a Stratagen MX3005P real time PCR machine (Stratagene, USA). Oligonucleotide primers are described as in Table (2).

Table 2: RT- PCR primers used in this study were supplied from Metabion (Germany).

RNA Virus	Gene	Primer/ probe sequence 5'-3'	Ref.
AI	Matrix	Sep1 AGATGAGTCTTCTAA CCGAGGTCG	Slomoka <i>et al.</i> 2007
		Sep 2 TGCAAAAACATCTTC AAGTCTCTG	
		SEPRO [FAM]TCAGGCCCC CTCAAAGCCGA [TAMRA]	
	H5	H5LH1 ACATATGACTAC CCACARTATTCA G	
		H5RH1 AGACCAGCT AYC ATGATTGC	
		H5PRO [FAM]TCWACA GTGGCGAGT TCCCTAGCA[TAMRA]	
	H9	H9F GGAAGAATTAATTATTATTGGTCGGTAC	Ben Shabat <i>et al.</i> 2010
		H9R GCCACCTTTTTCAGTCTGACATT	
		H9 Probe [FAM]AACCAGGCCAGACATTGCGAGTAAGATCC[TAMRA]	
ND	Vir	F+4839 TCCGGAGGATACAAGGGTCT	Wise <i>et al.</i> 2004
		F-4939 AGCTGTTGCAACCCCAAG	
		F+4894 [FAM]AAGCGTTTCTGTCTCCTTCCTCCA[TAMRA]	
	Matrix	M_4100 5_-AGTGATGTGCTCGGACCTTC-3_	
		M_4169 5_-[FAM]TTCTCTAGCAGTGGGACAGCCTGC[TAMRA]-3_	
		M_4220 5_-CCTGAGGAGAGGCATTTGCTA-3_	

Virus isolation:

The positive PCR samples were directed to virus isolation according to OIE 2008. Also virulence test evaluation for low pathogenic avian influenza virus (H9N2) and Newcastle virus. Depending on OIE 2012.

I: Pathogenicity of H9N2 virulence: protocol using 8 weeks SPF chicken. Virulence was evaluated for randomly selected two isolates:

II: Pathogenicity of NDV virulence: was done by injecting the virus in SPF day old chicks using ICPI.

Genotyping of isolates: sequencing for 20 positive isolates. It was performed by purification of PCR

products using QIAquick PCR Product extraction kit. (Qiagen, Valencia). The sequence reaction was done using Bigdye Terminator V3.1 cycle sequencing kit (Perkin-Elmer) and the sequence reaction was purified using Centrise spin column (ABI, USA). DNA sequences were obtained using Applied Biosystems 3130 genetic analyzer (ABI, USA). A phylogenetic tree was created by the MegAlign module of Lasergene DNASTar.

The sequencing method was done for HA gene of AI virus and for F gene of Newcastle virus.

RESULTS

A total 41 farms out of 195 examined are positive for AIV with a percentage of 21%, It appeared that 34 farm were positive for AI H9 with a percentage 83% and 7 farms were positive for AI H5 with a percentage of 17% Also 41 farms were positive for matrix gene of Newcastle virus with a percentage of 21%, Twenty four farms were positive for the virulent strain of Newcastle with a percentage of 58.5%.

Table 3: The results of PCR test for the examined Avian influenza and Newcastle viruses in relation to each governorate.

Gov.	No of examined farms	AI Matrix	The PCR results			
			AIH9	AI H5	ND	
					Matrix ND	vNDV
1- Faioum	51	15	14	1	12	7
2- Qalioubia	20	7	6	1	4	2
3- Dakahli	19	2	2	0	3	3
4- Ismalia	18	0	0	0	3	2
5- Benisueif	15	4	2	2	0	0
6- Domiatt	11	0	0	0	1	0
7- Luxor	12	2	1	1	1	0
8- Giza	9	0	0	0	2	2
9- Sharkia	10	3	1	2	3	1
10- Alexandria	8	3	3	0	7	4
11- Kafr el sheikh	5	1	1	0	3	2
12- Behira	5	1	1	0	1	0
13- Monefia	3	1	1	0	0	0
14- Gharbia	3	1	1	0	0	0
15- Minia	2	1	1	0	0	0
16- North Sinai	2	0	0	0	0	0
17- Aswan	1	0	0	0	0	0
18- Port said	1	0	0	0	1	1
Total		41	34	7	41	24
Incidence rate	195	21%	17.4%* 83%**	3.5%* 17%**	21%	12%* 58.5%**

Note: * the percent in relation to No of examined samples

** the percent in relation to the positive farms.

Virulence evaluation for different viruses:

I: evaluation of H9N2 virulence:

Virulence was evaluated for (2) isolates:

1-A/Chicken/Egypt/114940v/NLQP/2011/H9N2.

2-A/Chicken/Egypt/114922v/NLQP/2011/H9N2.

The results showed that the viruses that have been tested were low pathogenic.

II: virulence evaluation of isolated NDVs:

Using ICPI showed that, all evaluated viruses (NDV-EG-567f-2012) were virulent strains (ICPI=1.96).

Sequence analysis:

The genetic analysis for 4 strains of avian influenza viruses HA gene of the H5N1 type [A/chicken/Egypt/128s/2012, A/chicken/Egypt/12186F-9/2012, A/chicken/Egypt/12186F-12/2012 and A/chicken/Egypt/125s/2012] revealed that these viruses are highly pathogenic as well as tracking clade 2.2.1 virus which is endemic and widespread in Egypt since 2006 and no cases of Egyptian H5N1 variant strain which follows clade 2.2.1.1 and described in Table (4).


The genetic analysis of HA gene of 4 isolates of H9N2 revealed that they are of low pathogenic in the phylogenic tree they are tracking G1-lineage of Asian viruses which spread at Middle East and they are nearly similar to other Egyptian viruses from 2011 (Fig.2).

The genetic analysis of F gene of 12 selected isolates of NDV revealed that ten strains are virulent

genotype VII) and also the recent isolated strains from Middle East like Jordan (apmv1/chicken/Jordan/Jo 11/2011) and Israel (buzzard/Israel/714/2011) with a percentage of 99.98% while not similar to vaccinal strains as lasota with a percentage of 72% based on partial F gene. As well as 2 isolates of NDV were classified as avirulent that like vaccinal strain (Lasota – like NDV (lentogenic strain), (Fig.3).


Table 4: No. and type of isolates of each virus.

Virus	Genotype		No. of cases
H5N1	Classic A	Clade 2.2.1	4
NDV	Virulent	Vir VII	10
	Avirulent		2
H9N2	Low pathogenic	Asian G/ lineage	4




(Fig. 1): Phylogenetic analysis of HA gene of the Egyptian H5N1 isolates in comparison to other strains from Middle East region, Asian and other strains.

*Note: Black arrows refer to the sequenced strains in this study.



(Fig. 2): Phylogenetic analysis of HA gene of the Egyptian H9N2 isolates in comparison to other strains from Middle East region, Asian and other strains.

*Note : Black arrows refer to the sequenced strains in this study.



(Fig. 3): Phylogenetic analysis of F gene of the Egyptian NDV isolates in the present study 2012 in comparison to other strains.

*Note : Black arrows refer to the sequenced strains in this study.

DISCUSSION

Conventional microbiological methods are sensitive and specific, it takes considerable time (4 to 7 days) to reach a diagnosis; therefore, PCR-based detection and pathotyping assays have been developed to support/substitute for these traditional diagnostic tools and to replace animal experiments (Kant *et al.*, 1997; Aldous *et al.*, 2001; Creelan *et al.*, 2002; Pham *et al.*, 2005 and Wise *et al.*, 2004). Real-time RT-PCR can offer several advantages over standard RT-PCR, such as speed and lower risk of cross

contamination of clinical samples with previously amplified products. In addition, the detection of the amplicon with a sequence-specific probe further confirms the nature of the target, reducing false positivity (Schweiger *et al.*, 2000; Spackman *et al.*, 2002; Lee and Suarez, 2004; Stone *et al.*, 2004; Di Trani *et al.*, 2006 and Payungporn *et al.*, 2006). Standardized RT-PCR protocol can also be applied easily to other respiratory viruses. Our results for AI virus detection by RT-PCR confirm the endemic picture of AI H5 with a percentage of 17% from total examined farms and those strains belong to classic

strain in Egypt and these observation agree with results of Aly *et al.* (2008) which recorded that the HPAI virus of subtype H5N1 was first recorded in Egypt in 2006 and since then, the disease has become endemic and still causes a significant threat to the poultry industry and humans in Egypt. Detection of two subtypes of AI, HPAI H5N1 and LPAI-H9N2 is also detected in the present study with a percentage of 3.5%, 17.4% from total examined respectively, and this conclusion discussed by Shortridge *et al.*, 1998; Spackman *et al.*, 2002; Salzberg *et al.*, 2007 and Xu *et al.*, 2007 detected the presence of LPAI-H9N2 and the HPAI-H5N1 viruses have continuously co-circulated in domestic poultry in different areas. The high percent of AI H9 in this study is 83% from total positive AI cases it also reflects the low pathogenic nature of the virus which allow the silent spread of such viruses in commercial chickens and this results agree with that of (Nili and Asasi, 2002; Kim *et al.*, 2006; Aamir *et al.*, 2007; Naeem *et al.*, 2007 and Park *et al.*, 2011). Which discussed and recorded that phenomenon of silent spread of LP H9N2 in the Middle East region for several years indicated additional risk factor to the poultry industry. Although H9N2 viruses are characterized as low pathogenic avian influenza (LPAI) viruses, they may cause high morbidity and mortality Since avian H9N2 viruses are currently perceived to represent a significant threat to human health it is important to determine whether or not viruses of this subtype circulating in poultry in various parts of the world have the potential to infect people (Cameron *et al.*, 2000). To determine the genetic relationship of H5N1, H9N2 avian influenza viruses and ND virus currently prevalent in poultry in Egypt using Phylogenicity relationship analysis of the 4 selected strains of H5N1 are highly pathogenic as well as tracking subclade 2.2.1 classical strain and these results confirmed that recorded by Arafa *et al.* (2012a) that recorded the presence of HPAI H5N1, and that group is prevailing mainly in village poultry and had fewer mutations compared to the originally introduced virus in 2006 Since 2009, this group has started to be transmitted back to commercial sectors. On the other hand the Phylogenetic analysis of the HA gene showed that the 4 Egyptian isolates of H9N2 in our study were resemble that isolated from Egypt in 2011 the G1-like lineage similar to the circulating viruses in the Middle East with very close phylogeny to the Israeli viruses and these results also resemble the results obtained by Arafa *et al.* (2012b) when they recorded different strains of low pathogenic H9N2 in poultry sector. Currently the isolation of the causative virus using embryonated chicken eggs and its characterization by hemagglutination activity (HA)/ hemagglutination inhibition/in vivo pathogenicity assays is the standard protocol to confirm ND cases (Alexander 2004). The discrimination of a virulent (e.g., vaccine viruses) and virulent NDV strains is very important in regions

practicing vaccination as a control measure against ND, especially in the context of the differential diagnosis of avian influenza (Alexander 2004). In this study we use primers specific to amplify the F gene for Newcastle indeed similar primers were used by Kant *et al.* (1997). To amplify the F gene sequence specifically from virulent and non-virulent strains of NDV. In this surveillance 58.5% positive cases of Virulent strains of Newcastle disease virus were detected from total positive (NDV examined farms) this high percent is agree to that mentioned by Kim *et al.* (2013), that the Virulent NDV can cause destructive signs in chickens all over the world. Although the current vaccines are substantially effective, they do not completely prevent infection, virus shedding and disease. All NDV isolates belong to a single serotype. Consistent with this, currently used vaccines, such as strains B1 and LaSota, are known to protect against morbidity and mortality caused by NDV isolates in different parts of the world (Cho *et al.*, 2008). However, recent studies have suggested that NDV strains currently in circulation represent genotypes that differ from that of the vaccine strains (Miller *et al.*, 2010).

We conclude that the active surveillance is very important to detect the co-circulated viruses within the farms as well as evaluation of viruses and genotypic characterization help in the selection of the type of vaccine used.

REFERENCES

- Aamir, U.B.; Wernery, U.; Ilyushina, N. and Webster, R.G. (2007): Characterization of avian H9N2 influenza viruses from United Arab Emirates 2000–2003. *Journal of Virology*, 361, 45–55.
- Aldous, E.W.; Collins, M.S.; McGoldrick, A. and Alexander, D.J. (2001): Rapid pathotyping of Newcastle disease virus (NDV) with fluorogenic, probes in a PCR assay. *Vet. Microbiol.* 80:201–212.
- Alexander, D.J. (2004): Highly pathogenic avian influenza/Newcastle disease, p. 258–282. *In* OIE Manual of diagnostic tests and vaccines for terrestrial animals, 5th ed. Office International des Epizooties, Paris, France.
- Alexander, D.J. (2007): An overview of the epidemiology of avian influenza. *Vaccine*, 25, 5637–5644.
- Aly, M.M.; Arafa, A. and Hassan, M.K. (2008): Epidemiological findings of outbreaks of disease caused by highly pathogenic H5N1 avian influenza virus in poultry in Egypt during 2006. *Avian Dis.* 52:269–277.
- Arafa, A.; Suarez, D.; Kholosy, S.G.; Hassan, M.K.; Nasef, S.; Selim, A.; Dauphin, G.; Kim, M.; Yilma, J.; Swayne, D. and M.M. Aly, M.M. (2012a): Evolution of highly pathogenic avian influenza H5N1 viruses in Egypt indicating

- progressive adaptation. *Arch. Virol.* 157: 1931–1947.
- Arafa, A.; Hagag, N.M.; Yehia, N.; Zanaty, A.M.; Mahmoud M. Naguib, M.M. and Nasef, S. (2012b):* Effect of co-circulation of highly pathogenic avian influenza H5N1 subtype with low pathogenic H9N2 subtype on the spread of infections. *Avian Dis.*, 56:849-857.
- Ben Shabat, M.; Meir, R.; Haddas, R.; Lapin, E.; Shkoda, I. Raibstein, I.; Perk, S. and Davidson, I. (2010):* Development of a real-time TaqMan RT-PCR assay for the detection of H9N2 avian influenza viruses. *J Virol Methods.*; 168 (1-2): 72-7.
- Cameron, K.R.; Gregory, V.; Banks, J.; Brown, I.H.; Alexander, D.J.; Hay, A.J. and Lin, Y.P. (2000):* H9N2 subtype influenza A viruses in poultry in Pakistan are closely related to the H9N2 viruses responsible for human infection in Hong Kong. *Virology* 278, 36–41.
- Cho, S.H.; Kwon, H.J.; Kim, T.E.; Kim, J.H. and Yoo, H.S. (2008):* Characterization of a recombinant Newcastle disease vaccine strain. *Clin Vaccine Immunol* 15: 1572–1579. doi:10.1128/CVI.00156-08.PubMed: 18768673.
- Creelan, J.L.; Graham, D.A. and McCullough, S.J. (2002):* Detection and differentiation of pathogenicity of avian paramyxovirus serotype 1 from field cases with one-step reverse transcriptase-polymerase chain reaction. *Avian Pathol.* 31: 493–499.
- Di Trani, L.; Bedini, B.; Donatelli, I.; Campitelli, L.; Chiappini, B.; De Marco, M.A.; Delogu, M.; Buonavoglia, C. and Vaccari, G. (2006):* A sensitive one-step realtime PCR for detection of avian influenza viruses using a MGB probe and an internal positive control. *BMC Infect Dis* 6: 87.
- Duvvuri, V.R.; Duvvuri, B.; Cuff, W.R.; Wu, G.E. and Wu, J. (2009):* Role of positive selection pressure on the evolution of H5N1 hemagglutinin. *Genomics Proteomics Bioinformatics* (7): 47–56.
- El-Zoghby, E.F.; Arafa, A.; Hassan, M.K.; Aly, M.M.; Selim, A.; Kilany, W.H.; Selim, U.; Nasef, S.; Aggor, M.G.; Abdelwhab, E.M. and Hafez, M.H. (2012):* Isolation of H9N2 avian influenza virus from bobwhite quail (*Colinus virginianus*) in Egypt. *Arch Virol.* DOI10.1007/s00705-012-1269-z.
- Kant, A.; Koch, G.; Van Roozelaar, D.; Balk, F. and TerHuurne, A. (1997):* Differentiation of virulent and non-virulent strains of Newcastle disease virus within 24 hours by polymerase chain reaction. *Avian Pathol.* 26: 837–849.
- Kim, J.A.; Cho, S.H.; Kim, H.S. and Seo, S.H. (2006):* H9N2 influenza viruses isolated from poultry in Korean live bird markets continuously evolve and cause the severe clinical signs in layers. *Veterinary Microbiology*, 118,169–176.
- Kim, S.H.; Wanasen, N.; Paldurai, A.; Xiao, S.; Peter L.; Collins, P.L.; Siba, K. and Samal, S.K. (2013):* Newcastle Disease Virus Fusion Protein Is the Major Contributor to Protective Immunity of Genotype-Matched Vaccine., Vol8 (8): e74022.
- Lee, M.S.; Chang, P.C.; Shien, J.H.; Cheng, M.C. and Shieh, H.K. (2001):* Identification and subtyping of avian influenza viruses by reverse transcription-PCR. *J. Virol Methods* 97, 13–22.
- Lee, C.W. and Suarez, D.L. (2004):* Application of real-time RT-PCR for the quantitation and competitive replication study of H5 and H7 subtype avian influenza virus. *J. Virol Methods* 119: 151–158.
- Miller, P.J.; Decanini, E.L. and Afonso, C.L. (2010):* Newcastle disease: evolution of genotypes and the related diagnostic challenges. *Infect Genet Evol* 10: 26-35. doi: 10.1016/j.meegid.2009.09.012. PubMed: 19800028.
- Mohamed, M.H.; Kumar, S.; Paldurai, A.; Megahed, M.M.; Ghanem, I.A.; Lebda, M.A. and Samal, S.K. (2009):* Complete genome sequence of a virulent Newcastle disease virus isolated from an outbreak in chickens in Egypt. *Virus Genes*, 39: 234-237.
- Munch, M.; Nielsen, L.P.; Handberg, K.J. and Jorgensen, P.H. (2001):* Detection and subtyping (H5 and H7) of avian type A influenza virus by reverse transcription-PCR and PCR-ELISA. *Arch Virol* 146, 87–97.
- Naeem, K.; Siddique, N.; Ayaz, M. and Jalalee, M.A. (2007):* Avian influenza in Pakistan: outbreaks of low- and high-pathogenicity avian influenza in Pakistan during 2003-2006. *Avian Diseases*, 51(1 Suppl), 189-193.
- Nili, H. and Asasi, K. (2002):* Natural cases and an experimental study of H9N2 avian influenza in commercial broiler chickens of Iran. *Avian Pathology*, 31, 247–252.
- OIE (2008):* Manual of Diagnostic tests and Vaccines for Terrestrial Animals,
- OIE (2012):* Manual of Diagnostic tests and Vaccines for Terrestrial Animals, Chapter 2.3.4.
- Park, K.J.; Kwon, H.I.; Song, M.S.; Pascua, P.N.; Baek, Y.H.; Lee, J.H.; Jang, H.L.; Lim, J.Y.; Mo, I.P.; Moon, H.J.; Kim, C.J. and Choi, Y.K. (2011):* Rapid evolution of low pathogenic H9N2 avian influenza viruses following poultry vaccination programmes. *J. Gen. Virol.* 92, 36–50.
- Payungporn, S.; Chutinimitkul, S.; Chaisingh, A.; Damrong wantanapokin, S.; Buranathai, C.; Amonsin, A.; Theamboonlers, A. and Poovorawan, Y. (2006):* Single step multiplex real-time RT-PCR for H5N1 influenza A virus detection. *J Virol Methods* 131:143–147.
- Perk, S.; Golender, N.; Banet, C.; Shihmanter, E.; Pirak, M.; Tendler, Y.; Lipkind, M. and*

- Panish, A. (2009):* Phylogenetic analysis of hemagglutinin, neuraminidase, and nucleoprotein genes of H9N2 avian influenza viruses isolated in Israel during the 2000–2005 epizootic. *Comparative Immunology, Microbiology and Infectious Diseases*. 32, 221–238.
- Pham, H.M.; Konnai, S.; Usui, T.; Chang, K.S.; Murata, S.; Mase, K.; Ohashi, M. and Onuma, M. (2005):* Rapid detection and differentiation of Newcastle disease virus by real-time PCR with melting-curve analysis. *Arch. Virol.* 150: 2429–2438.
- Salzberg, S.L.; Kingsford, C.; Cattoli, G.; Spiro, D.J.; Janies, D.A.; Aly, M.M.; I. Brown, H.; Couacy-Hymann, E.; De Mia, G.M.; Dung do, H.; Guercio, A.; Joannis, T.; Maken Ali, A.S.; Osmani, A.; Padalino, I.; Saad, M.D.; Savic, V.; Sengamalay, N.A.; Yingst, S.; Zaborsky, J.; Zorman-Rojs, O.; Ghedin, E. and Capua, I. (2007):* Genome analysis linking recent European and African influenza (H5N1) viruses. *Emerg. Infect. Dis.* 13: 713–718.
- Schweiger, B.; Zadow, I.; Heckler, R.; Timm, H. and Pauli, G. (2000):* Application of a fluorogenic PCR assay for typing and subtyping of influenza viruses in respiratory samples. *J Clin Microbiol* 38: 1552–1558.
- Shortridge, K.F.; Zhou, N.N.; Guan, Y.; Gao, P.; Ito, T.; Kawaoka, Y.; Kodihalli, S.; Krauss, S.; Markwell, D.; Murti, K.G.; Norwood, M.; Senne, D.; Sims, L.; Takada, A. and Webster, R.G. (1998):* Characterization of avian H5N1 influenza viruses from poultry in Hong Kong. *Virology* 252, 331–342.
- Slomka, M.J.; Pavlidis, T.; Banks, J.; Shell, W.; McNally, A.; Essen, S. and Brown, I.H. (2007):* Validated H5 Eurasian real-time reverse transcriptase–polymerase chain reaction and its application in H5N1 outbreaks in 2005–2006. *Avian Dis* 51: 373–377.
- Spackman, E.; Senne, D.A.; Myers, T.J.; Bulaga, L.L.; Garber, L.P.; Perdue, M.L.; Lohman, K.; Daum, L.T. and Suarez, D.L. (2002):* Development of a real-time reverse transcriptase PCR assay for type A influenza virus and the avian H5 and H7 hemagglutinin subtypes. *J Clin Microbiol* 40: 3256–3260.
- Starick, E.; Romer-Oberdorfer, A. and Werner, O. (2000):* Type- and subtype-specific RT-PCR assays for avian influenza A viruses (AIV). *J. Vet Med B Infect Dis Vet Public Health* 47, 295–301.
- Stone, B.; Burrows, J.; Schepetiuk, S.; Higgins, G.; Hampson, A.; Shaw, R. and Kok, T. (2004):* Rapid detection and simultaneous subtype differentiation of influenza A viruses by real time PCR. *J Virol Methods* 117: 103–112.
- Wise, M.G.; Suarez, D.L.; Seal, B.S.; Pedersen, J.C.; Senne, D.A.; King, D.J.; Kapczynski, D.R. and Spackman, E. (2004):* Development of a real-time reverse-transcription PCR for detection of Newcastle disease virus RNA in clinical samples. *J. Clin. Microbiol.* 42: 329–338.
- Xu, K.M.; Smith, G.J.; Bahl, J.; Duan, L.; Tai, H.; Vijaykrishna, D.; Wang, J.; Zhang, J.X.; Li, K.S. and Fan, X.H. (2007):* The genesis and evolution of H9N2 influenza viruses in poultry from southern China, 2000 to 2005. *J. Virol* 2007, 81:10389–10401.

استبيان وتوصيف فيروس انفلوانزا الطيور وفيروس النيوكاسل في قطاع التسمين والبيض في مصر

أزهار جابر على شلبي ، أحمد محمد عبد الرحمن عرفان ، فاطمة عامر عبد الرحيم ، عبد الله عبد الظاهر سليم ،
محمد حسن عبد الله ، شيرين جلال خلوصي ، سعاد عبد العزيز عبد الونيس

Email: azhar_gaber@yahoo.com

نظرا لانتشار فيروس الانفلوانزا H5N1 في قطاع الدواجن في مصر منذ مدة فقد اوضح الاستبيان النشاط في ١٩٥ مزرعة تسمين وبيض لثمانية عشر محافظة مصرية ٣٤، ٧، ٢٤ مزرعة ايجابية ل H9N2 و H5N1 والنيوكاسل الضاري على الترتيب ووضح التحليل الجيني لجين الهيماجلوتينين لفيروس انفلوانزا الطيور للعترة H5N1 أظهر أنها عترات عالية الضراوة كلاسيكية من الفصيلة ٢.٢.١ والتي استوطنت وانتشرت في مصر منذ ٢٠٠٦. لكن عترات ال H9N2 كانت منخفضة الضراوة وتنتمي ل G1-lineage من الفيروسات الأسبوية والتي تنتشر في الشرق الأوسط. أيضا تحليل التتابع النيوكليتيدي لجين F لإثنى عشر معزولة نيوكاسل أظهر أنهم شديدي الضراوة ماعدا معزولتين كانتا منخفضتي الضراوة وتشابه عترة التحصين لاسوتا. تم تقييم الضراوة لبعض معزولات النيوكاسل من خلال معامل الضراوة بالحقن المخي كما حدث مع العترة (NDV-EG-567f-2012) والتي ظهرت أنها عالية الضراوة وكانت قيمة معامل ضراوة الحقن المخي ١.٩٦. ومن هذه الدراسة تم إثبات تواجد فيروس الانفلوانزا من العترة H5N1 والعترة H9N2 منخفضة الضراوة في الطيور في نفس الوقت في مصر.