

## STUDY THE PREVALENCE OF PSYCHROTROPHIC FOOD BORNE BACTERIA IN *TILAPIA NILOTICA* IN DAKAHLIA MARKETS AND ITS EFFECT ON FISH QUALITY

SHEREEN S. MOUSTAFA; HATEM F.A. EL-DOSOKY and AZA. E.A. HASSAN  
Animal Health Research Institute, Mansoura Branch

---

### ABSTRACT

Received at 27/2/2013

Accepted: 11/4/2013

One hundred samples of *Tilapia nilotica* were collected from different markets in Dakahlia Governorate to evaluate their bacteriological quality by determine Psychrotrophic counts. The effect of these foodborne bacteria on chemical quality of fish and consumer health were studied. The bacteriological examination revealed that the mean value of total Psychrotrophic count, total *Pseudomonas* and total *Bacillus cereus* count in *Tilapia nilotica* samples were  $3.1 \times 10^2 \pm 1.8 \times 10^2$  cfu/ gram,  $5.1 \times 10^3 \pm 1.4 \times 10^2$  cfu/ gram and  $4.0 \times 10^3 \pm 2.6 \times 10^2$  cfu/gram respectively. Biochemical tests as Histamine (H), Thiobarbituric acid (TBA), Total volatile basic amines (TVBN) were measured to determine the quality of examined fish and their effect on health of consumers. The obtained results revealed that, the minimal and maximal histamine(H) content were 5.7 and 24.6 mg% respectively, 10.2 & 25.1 mg/100gm for minimal and maximal (TVBN) respectively and 0.534 & 0.988 mgDK/Kg for minimal and maximal (TBA) content in examined fish samples

---

**Key words:** Fish -*Tilapia nilotica* -*Bacillus cereus* -*Pseudomonas* - Psychrotrophic

---

### INTRODUCTION

Fish and fish products were incriminated as a cause of food poisoning, food intoxication as well as many infectious diseases (Lawsan, 1970). The microflora on fish is dominated by psychrotrophic Gram-negative rod shaped bacteria belonging to the genera *Pseudomonas*, *Moraxella*, *Acinetobacter*, *Shewanella* and *Flavobacterium*. Members of the *Vibrionaceae* (*Vibrio* and *Photobacterium*) and the *Aeromonadaceae* (*Aeromonas* spp.) are also common aquatic bacteria and typical of the fish flora. Gram-positive organisms as *Bacillus*, *Micrococcus*, *Clostridium*, *Lactobacillus* and coryneforms can also be found in varying proportions, but in general, Gram-negative bacteria dominate the microflora. Some psychrotrophic pathogens can grow in refrigerated food with little or no obvious change of sensory characteristics (Berrang *et al.*, 1989) *Pseudomonas* species and *Bacillus cereus* are the most predominant psychrotrophic microorganisms and their presence in food creates a great risk as they lead to food poisoning and / or spoilage of food products (Jay, 2000). Also *Bacillus cereus* causes health hazard to consumers as they cause spoilage and deterioration to meat and meat products and responsible for food poisoning outbreaks (Parry *et al.*, 1993). *Bacillus cereus* can give rise to two distinct forms of food borne diseases, the emetic form and diarrheal form, the emetic form believed to be associated with an emetic toxin performed in food

while diarrheal form is caused by an enterotoxin (Altayer and Sutherland, 2006). From the food safety point of view, proteins as normal constituent of fish are exposed to degradation by microbial enzymes activity specially proteases (Gill, 1990), Decarboxylation of amino acids histidine is encountered among several species of bacteria including *Bacillus cereus* and *Pseudomonas* the significance important of histamine in fish has been discussed by several authors (Vidal-carroa *et al.*, 1990 and Hod & Khalafalla, 1993). Encountered among several species of bacteria from another point of view we study the effect of psychrotrophic bacteria on the chemical changes of fish, it is well known that both enzymatic and microbiological activity are greatly influenced by temperature. So the aim of this study is the detection of the most common psychrotrophic food borne pathogens in *Tilapia nilotica* fish samples collected from different markets in Dakahlia Governorate and evaluate their effect on fish quality.

### MATERIALS and METHODS

#### 1 – Collection of samples

- A total of 100 random fish samples of *Tilapia nilotica* were collected from fish markets in Dakahlia Governorate.
- Collected samples were transferred in ice box immediately to the laboratory under complete aseptic precautions without delay for bacteriological examination and quality tests.

**2- Preparation of samples for bacteriological examination:-**

- 10 gram of each fish and 90 ml of sterile peptone water were thoroughly mixed by sterile blender for approximately 2.5 min., from which ten fold serial dilutions were prepared.

- The prepared samples were subjected to the following examination:-

**a- Determination of psychrotrophic bacterial count (ISO, 2004);** using surface spread technique 0.1 ml from each dilution was transferred and evenly spread over a dry surface of previously prepared sterile plate count agar media, allowed to dry for 15 min., inoculated plates divided into two groups each containing one plate from each dilution, the first group incubated at 7° for 10 days and second incubated at 25° for 24 hr.

- Average number of colonies per gram was determined and the psychrotrophic count / gram was calculated and recorded.

**b- Enumeration and identification of Pseudomonas species (ISO, 2004)**

- 0.1 ml of each dilution was separately inoculated into duplicated Petri-dishes of Pseudomonas selctive

agar media supplemented with glycerol. The inoculated plates were incubated at 25° c for 48hr after which all developed colonies with greenish yellow pigment were enumerated and the average count / gram was calculated and recorded.

**c- Enumeration and identification of Bacillus cereus (ISO, 1987)**

- The spreading technique was applied on the surface of Bacillus cereus selective agar media which incubated at 30° c for 24hr, then count was recorded. Suspected Colonies were identified by microscopical and biochemical examination.

**3- CHEMICAL EXAMINATION OF FISH**

**a- Quantitative Assessment of Histamine:** was done by using thin layer chromatography method (TLC), (INFO SAMAK, 1989)

**b- Determination of total volatile basic nitrogen (TVBN) mg/100gm:** It was done according to FAO (1980).

**c- Quantitative Assessment of Thiobarbituric acid (TBA) MD/kg:-** According to Pikul *et al.* (1983)

**RESULTS**

**Table 1:** Total Psychrotrophic count, Pseudomonas and Bacillus cereus counts of the examined Tilapia nilotica samples (n= 100)

Bacterial count	Total psychrotrophic	<i>pseudomonas</i>	<i>Bacillus cereus</i>
<i>Minimum</i>	2.0 ×10 <sup>3</sup>	8.2×10 <sup>2</sup>	7.0 × 10 <sup>2</sup>
<i>Maximal</i>	4.2 ×10 <sup>4</sup>	2.1×10 <sup>4</sup>	8.2 × 10 <sup>3</sup>
<i>Mean</i>	3.1×10 <sup>2</sup>	5.1×10 <sup>3</sup>	4.0× 10 <sup>3</sup>
<i>Standard error</i>	1.8 ×10 <sup>2</sup>	1.4× 10 <sup>2</sup>	2.6 × 10 <sup>2</sup>

**Table 2:** Frequency distribution of examined Tilapia nilotica samples based on their total Psychrotrophic & Pseudomonas and Bacillus cereus count. (n= 100):

Count Range	Total Psychrotrophic		<i>Pseudomonas</i>		<i>Bacillus cereus</i>	
	No.	%	No.	%	No.	%
<10 <sup>2</sup>	45	45	40	40	38	38
10 <sup>2</sup> < 10 <sup>3</sup>	7	7	16	16	22	22
10 <sup>3</sup> <10 <sup>4</sup>	13	13	8	8	40	40
10 <sup>4</sup> < 10 <sup>5</sup>	35	35	36	36	0	0

Negative < 10<sup>2</sup> (according to Amal F.A & Shaboury, 2009)

**Table 3:** Results of biochemical examination of fish samples:-  
N. of samples = 25 of each

Biochemical test	Minimum	Maximum	Mean ±S.e
<b>Histamine</b> Mg%	5.7	24.6	11.2±1.3
<b>TVB- N</b> mg/100gm	10.2	25.1	16.3±3.2
<b>TBA</b> mg D M/Kg	0.534	0.988	0.701±0.04

**Table 4:** Incidence of Histamine level Histamine, TVB-N and TBA in *T. nilotica* fish samples in relation to the level pointed by Egyptian standard:

Kind of fish	Biochemical test	No of samples	Permissible limit	Lower than permissible limit		No higher than permissible limit	
				No.	%	No.	%
<i>Tilapia nilotica</i>	-Histamine mg%	25	10mg/100 gmuscle	21	84.0	4	16.0
	-TVB- N mg/100gm	25	25mg/100gm	22	88.0	3	12.0
	- TBA mg D M/Kg	25	4.0 mg DM/Kg	25	100	0	0.0

## DISCUSSION

Fish are regarded as being most popular and more perishable than other high protein foods. The flesh of healthy fish is considered bacteriologically sterile, However they are sometimes contaminated with bacterial pathogens and thus can inflict heavy losses in fish all over the world. Fish and fish products were incriminated as a cause of food poisoning, food intoxication as well as many infectious diseases (Lawsan, 1970). The Psychrotrophic bacteria have been received an increased attention by several investigators during recent years due to modern developments in fish production which results in fish must be held for longer period at low temperature before transportation, processing, manufacture or consumption. Psychrotrophes are those organisms that grow well at or below 7° c and have their optimum between 20°c - 30°c. Some psychrotrophic pathogens can grow in refrigerated food with little or no obvious change of sensory characteristics (Berrang *et al.*, 1989) *Pseudomonas* species and *Bacillus cereus* are the most predominant psychrotrophic micro organisms and their presence in food creates a great risk as they lead to food poisoning and / or spoilage of food products (Jay, 2000). The results in table (1) revealed that the total Psychrotrophic count of examined *Tilapia nilotica* samples ranged from  $2.0 \times 10^3$  as a minimum count to  $4.2 \times 10^4$  as a maximal count with a mean  $3.1 \times 10^2 \pm 1.8 \times 10^2$  SE, variable results were reported by (Mousa and Mahmoud, 1997), who recorded that the

mean values of total Psychrotrophic count of examined *tilapia nilotica* was  $0.56 \times 10^2 \pm 0.028 \times 10^2$  /g. While (Mahmoud, 1994). Recorded that the Mean value was  $5.84 \times 10^3 \pm 0.090$ . (Amal F.A El. Shaboury, 2009). mentioned that the total Psychrotrophic count of examined *Tilapia nilotica* samples ranged from  $1.7 \times 10^3$  to  $3 \times 10^4$  with a mean value of  $3.2 \times 10^2 \pm 1.3 \times 10^2$ . Results in table (2) mentioned that the frequency distributions of total Psychrotrophic count of examined *Tilapia nilotica* samples were 45 % out of the examined samples considered negative ( $< 10^2$ ). 7% of samples lied between  $10^2$  to  $< 10^3$ , 13 % lied between  $10^3$  to  $< 10^4$  while the majority of positive samples (35%) lied between  $10^4$  to  $< 10^5$ . The obtained results agreed with that mentioned by (Hayes, 1992), and (Amal F.A & El-Shaboury 2009) as they recorded that the majority of examined *Tilapia nilotica* samples lied between  $10^4$  to  $10^5$ . (Lawsan, 1970), stated that fish, fish products were incriminated as a cause of food poisoning, food intoxication as well as many infectious diseases. *Pseudomonas* species and *Bacillus cereus* are the most predominant Psychrotrophic bacteria which their presence in food creates a great risk as they lead to food poisoning and / or spoilage of food products (Jay, 2000). *Pseudomonas* species was isolated from fresh water fish by Gram 1993, (Mousa and Mahmoud, 1997), in this study results in table (1) mentioned that the minimum *Pseudomonas* species count of examined *Tilapia nilotica* samples was  $8.2 \times 10^2$  & maximal count was  $2.1 \times 10^4$  with means count of  $5.1 \times 10^3 \pm 1.1 \times 10^2$ . Lower *Pseudomonas* count

was recorded by (Lamada – Hanan, 1999), but this above results agreed with that recorded by (Amal F.A and El-Shaboury, 2009) as they recorded that the mean value of pseudomonas count in the examined Tilapia nilotica samples was  $4.5 \times 10^3$ . The frequency distribution of examined tilapia nilotica samples based on their pseudomonas count was recorded in table (2) and revealed that the majority of samples lied between  $10^4 < 10^5$  with the percentage of 36% from total examined tilapia nilotica samples, a same results were recorded by (Hassan 1991 and Amal F.A & El- Shaboury 2009). *Bacillus cereus* was widely distributed in nature, water, soil, air and can be isolated from a wide variety of foods, (Parry *et al.*, 1993) mentioned that *Bacillus cereus* cause health to consumers and also responsible for food poisoning outbreaks. The results in table (1) illustrated that the minimal *Bacillus cereus* count of examined Tilapia nilotica samples was  $7.0 \times 10^2$ , maximal was  $8.2 \times 10^3$  with mean value  $4.0 \times 10^3 \pm 2.6 \times 10^2$ . The frequency distribution of *Bacillus cereus* count in tilapia nilotica samples was recorded in table (2) which revealed that the majority of samples lied between  $10^3 < 10^4$ , a resembling results were mentioned by (Amal F.A & El-Shaboury, 2009). In this study the quality outlines of examined Tilapia nilotica were detected by determining the effect of Psychrotrophic bacteria on biochemical characters of examined fish, a quantitative assessment of Histamine (H), Total volatile basic nitrogen (TVBN) and Thiobarbituric acid (TBA) was made to determine the effect of *Pseudomonas* species and *Bacillus cereus* on fish quality. Results in table (3) revealed that minimal and maximal level of histamine were 5.7 and 24.6 mg% with  $11.2 \pm 1.3$ SE mean value respectively, these were nearly in agreement with results registered by (Park *et al.*, 1980), (Vidal –Caroia *et al.*, 1990); (Hoda, H.A. and Khalafala, 1993), as they recorded that histamine level in naturally spoiled fish ranged from 22 mg/Kg up to 25 mg/kg. (Hosseini *et al.*, 2009) mentioned that histamine amounts depend on production date of samples. Table 3 showed that minimal and maximal values of TVBN were 10.2 & 25.1 mg/100gm respectively with a mean value  $16.3 \pm 3.2$  mg/100gm these results with in agreement with (Acuff *et al.*, 1984). Increasing of TVBN during storage indicated possible spoilage of fish, such increase may be attributed to the production of volatile basic compound such as ammonia as reported by (Putro *et al.*, 1985; Galli *et al.*, 1993). Also table (3) revealed that the minimal and maximal thiobarbituric acid (TBA) values of examined fish samples were 0.534 mg MD/Kg and 0.988 mg MD/Kg with a mean value  $0.701 \pm 0.04$ mg DM/Kg and these results were in agreement with (Undeland and Lingnert 1999). However these results were lower than (Sohad *et al.*, 2008) who recorded an increase in thiobarbituric acid value (TBA). From all above results fresh fish should be consumed as soon as possible and surely within 3 days from purchased

as recorded by (Ahmed and Yassien 2000), because of longer storage may result in rapid rise in histamine content, volatile nitrogen basic compound and thiobarbituric acid. Also growth of Psychrotrophic bacteria in fish has become a significant problem due to wide spread use of refrigerated storage of fish, that cause a potential threat to consumers health. According to the results recorded in table (4), Histamine, TVBN and TBA, were higher than the permissible limits obtained by Egyptian standard (2005) with percentages 16.0, 12.0 and 0.0. respectively.

## REFERENCES

- Acuff, G.; Izat, A.L. and Finne, G. (1984): Microbial flora of reared tilapia held on ice. J. Food. Prot., 47: 778–780.
- Ahmed, A.M. and Yassien, M.A (2000): Level of histamine-forming bacteria in fish from Ismailia markets with records of scombroid poisoning in children. S.C.V. M.J., III (1).
- Altayer, M. and Sutherland, A.D. (2006): *Bacillus cereus* is a common in the environment but emetic toxin producing isolates are rare. J. Appl. Microbiology, 100 (1): 7-14.
- Amal, F.A. Mansour and EL-Shaboury, F.A. (2009): Prevalence of psychrotrophic foodborne pathogens in fish in Alexandria markets. Assiut Vet. Med. J. Vol. 55. No. 121 April 2009.
- A.P.H.A (American Public Health Association) (1984): compendium of Methods for Microbiological Examination of foods. 3<sup>ed</sup> Ed. Washington, D.C.
- Berrang, M.E.; Brachett, R.E. and Beuchat, L.R. (1989): Groth of listeria monocytogenes on fresh vegetables stored under a controlled atmosphere. J. Food prot. (52): 702-705.
- Campton, R. (1981): In Campton Encyclopedea vol. 5 pp. 601. Publisher University of Chicago VSA.
- FAO (1980): Manual of Food Quality Control. IV. Microbiological analysis. FAO United Nations, Rome. Tech. Res. Ser.
- Galli, A.I.; Franzetti, S.; Carelli, L.; Piergiovanni, I. and Fava, P. (1993): Microbiological quality and shelf life of chilled cod fillets in Vacuum – skin and modified atmosphere packaging. Pack. Technol. Sci., 6: 147–154.
- Hassan, M.T. (1991): Microbiological status of frozen dressed fish. M.V.SC Thesis Fac. Vet. Med., Moshtohor, Zagazig Univ. – Banha.
- Hayes, P.R. (1992): Food Microbiology and Hygiene. Elsevier science publishers (L.T.d).
- Hoda, H. Awad and Khalafalla, F. (1993): Histamine level in imported scombroid fishes. Vet. Med. J., Giza 41, 3: 67– 71.
- Hosseini, H.; Dolatabadi, R.; Shekarchi, M.; Keshavarz, A.; Eskanadari, S. and Pirali-Hamedani, M. (2009): Evaluation of histamine

- in canned tuna fish post market samples in Iran using Elisa Asian J. of Chemistry. 21, 8: 6429–6434.
- ICSMF (International commission on Microbiological specification for foods) (1978):* Their significance and methods of enumeration. 2<sup>nd</sup> Ed. Univ. of Toronto press. Toronto - buffalo, Canda.
- INFO SAMAK/UNDP/FAO (1989):* Training course on fish processing technology and quality control. Cairo, Egypt, 28 January -15 February.
- ISO (1987):* Microbiology general guidance for enumeration of Bacillus cereus. ISO 7932. Geneva, Switzerlan.
- ISO 7932 (2004):* Microbiology of food and animal feeding stuffs, Horizontal methods for enumeration of presumptive Bacillus cereus – colony count technique at 30 dgree c.
- Jay M.T. (2000):* Modern Food microbiology. Sixth Ed. Gaithersburg. Maryland.
- Lamada–Hana, M.O. (1999):* Chemical and sensory changes associated with microbial flora of Miditerranean boque (Boops –boops) stored aerobically at 0, 3, 7 and 10°. Appl. Environ. Microb. 65 (2): 698.
- Lawsan, T.B. (1970):* Some aspects of fish inspection and public health Vet. Reas., 87: 52.
- Mahmoud, Y.E. (1994):* Studies on frozen fish. Ph. D. thesis, Fac. Vet. Med. Moshtoher, Zagazig University. Benha.
- Mousa, M.M. and Mahmoud, Y.E. (1997):* Hygenic quality of newly caught fresh fish. Assuit. Vet. Med. J. 219-223.
- Park, Y.H.; Kim, D.S. and Kim, S.B. (1980):* Changes in histamine content in muscle of dark fleshed fish during storing and processing. Bull. Korean fish Soc., 13: 15-22.
- Parry, J.M.; Turnbull, P.C.B. and Gibson, J.R. (1993):* Colour Atlas of Bacillus species, Wolf Medical Publication Ltd. London.
- Pikul, J.; Leszczynski, D.E. and Kummerow, F. (1983):* Elimination of sample autoxidation by butylatedhydroxyl toluene. J. Agric. Food Chem. 31: 1338.
- Putro, S.; Saleh, M. and Bandolutmoto, B.S. (1985):* Storage life of rabbit during icing. FAO Fisheries. Rep., Np 317 (suppl.), 54: 61.
- Sohad, H.E.; EL. Leboudi, Nahed; Salem, I.F. and Heikal, G.I. (2008):* Bacterial and chemical evaluation of some imported frozen fish in Kafer–El-Sheikh Governorate markets. J. Egypt. Vet. Med. Assoc. 68, 3: 119-128.
- Undeland, I. and Lingnert, H. (1999):* Lipid oxidation in fillets of herring (Glupea harengus) during frozen storage. J. of Agr. and Food Chemi., 47: 275-279.
- Vidal-Carao, M.C.; Maria, T.V. and Font, A.M. (1990):* Spectrofluorometric determinate of histamine in fish and meat products. J. Assoc. of Anal. Chem., 73, 4: 565–567.

### دراسة مدى تواجد البكتريا الممرضة المحبة للبرودة فى اسماك البلطى النيلية فى اسواق محافظة الدقهلية وتأثيرها على جودة الاسماك

شيرين سامى مصطفى ، حاتم فتحى احمد الدسوقي ، عزه السيد حسان

تم جمع عدد 100 عينة من اسماك البلطى النيلية من اسواق محافظة الدقهلية وذلك لتقييمها من الناحية الصحية ودراسة مدى تواجد البكتيريا المحبة للبرودة وبيان مدى تأثيرها على جودة الاسماك. وقد دلت النتائج على ان متوسط العد الكلى للميكروبات المحبة للبرودة  $3.1 \times 10^2$ /gram بينما كان متوسط العد الكلى لميكروب السودوموناس  $5.1 \times 10^3$ /gram ومتوسط العد الكلى لميكروب الباسيليس سيرس  $4.0 \times 10^3$ /gram. تحتوى لحوم الاسماك على نسبة عالية من الاحماض الدهنية بالاضافة الى الاحماض الامينية والتي تتعرض للتحلل نتيجة للنشاط البكتيرى وقد تم اجراء اختبارات كيميائية عن طريق تقدير نسبة الهيستامين – القواعد النيتروجينية الكلية - حامض الثيوباربتيوريك للكشف عن جودة الاسماك وكانت النسب كالاتى: الحد الادنى للهيستامين كان 5.7 مجم % والحد الاقصى 25.1 مجم % اما بالنسبة للقواعد النيتروجينية الكلية وحامض الثيوباربتيوريك فقد كانت نسب الحد الادنى والحد الاقصى  $10.2\text{mg}/100\text{gm}$  &  $25.1\text{mg}/100\text{gm}$  للقواعد النيتروجينية و  $0.534\text{ mgDK}/\text{Kg}$  &  $0.988\text{mgDK}/\text{Kg}$  لحامض الثيوباربتيوريك.