

Department of Food Hygiene,  
Faculty of Veterinary Medicine,  
Assiut University, Assiut, Egypt.

## **INVESTIGATION OF MILK AND SOME DAIRY PRODUCTS FOR FECAL POLLUTION INDICATORS** (With 6 Tables)

By

**ENAS EL-PRINCE; MOHAMMED SAYED\*; MAHMOUD  
FARGHALY\*\* and AMIRA M. ABDEL-RAHMAN\*\***

\*Corresponding author (dr.mohammedsayed@yahoo.com)

\*\*Animal Health Research Institute, Assiut Regional Laboratory, Egypt.

(Received at 29/8/2010)

### **فحص اللبن وبعض منتجات الألبان لمؤشرات التلوث البرازية**

**إيناس البرنس ، محمد سيد ، محمود فرغلي ، أميرة محمد عبد الرحمن**

تم إجراء هذه الدراسة للتعرف على مدى تلوث الألبان وبعض منتجاتها بهذه الميكروبات. لذلك تم جمع 180 عينة عشوائية من اللبن الخام، الجبن القريش، الجبن الدماطي، الجبن المطبوخ، الأيس كريم و الزبادي بواقع 30 عينة من كل نوع من الأسواق المختلفة بمدينة أسيوط. وقد أسفرت النتائج على أن جميع عينات اللبن الخام والجبن القريش كانت ملوثة بالميكروبات القولونية بينما تواجدت بنسبة 23.3، 16.7، 93.3 و 46.7% في عينات الجبن الدماطي والجبن المطبوخ والأيس كريم والزبادي على التوالي. وكانت 60، 83.3، 6.7، 6.7، 56.7 و 36.7% من العينات المفحوصة على الترتيب ملوثة بالميكروبات القولونية البرازية. أما ميكروب الإشيريشياكولاي فقد تم عزله بنسبة 3.3، 3.3، 33.3، 3.3، 6.7 و 20% من العينات على الترتيب. كما أتضح أن 83.3، 83.3، 26.7، 20، 70 و 40% من العينات كانت تحتوي على ميكروب الإنتيروكوكاي. وقد وجد الكلوستريديم بيفرنجينز فقط في عينات الجبن القريش بنسبة 3.3%. ويمكن أن نستخلص من نتائج الدراسة أن النسبة العالية من الإنتيروكوكاي تجعلها المفضلة للكشف عنها كمؤشر للتلوث البرازي.

**الكلمات الكاشفة:** اللبن، منتجات الألبان، مؤشرات التلوث البرازية، الإشيريشياكولاي، الإنتيروكوكاي، الكلوستريديم بيفرنجينز.

## **SUMMARY**

The goal of this study was to investigate milk and some dairy products for fecal pollution indicators. Therefore, 180 random samples of raw

milk, Kareish cheese, Domiati cheese, processed cheese, ice cream and yoghurt (30 each) were collected from Assiut city. All the examined raw milk and Kareish cheese samples were contaminated with coliforms, and their percentages in Domiati cheese, processed cheese, ice cream and yoghurt samples were 23.3, 16.7, 93.3 and 46.7%, respectively, while, 60, 83.3, 6.7, 6.7, 56.7 and 36.7% of the examined samples were contaminated with fecal coliforms, respectively. 3.3, 33.3, 3.3, 3.3, 6.7 and 20% of the examined samples were contaminated with *E. coli*, respectively. The incidences of *Enterococci* were 83.3, 83.3, 26.7, 20, 70 and 40%, respectively, while *Cl. perfringens* was detected only in Kareish cheese samples in a percentage of 3.3%. It can be concluded that the high level of *Enterococci* in the examined milk and milk products samples declared that *Enterococci* is preferable as an indicator for fecal pollution.

**Key words:** Milk, Milk products, Fecal pollution indicators, *E. coli*, *Enterococci*, *Cl. perfringens*.

## INTRODUCTION

Milk is a food of complex biochemical composition and high water activity serving it as an excellent culture medium for the growth and multiplication of many kinds of microorganisms (Oliver *et al.*, 2005). Among these microorganisms a group of bacteria called enteric pathogens which represent one of the most important bacterial groups in food. Members belonging to this group can survive and multiply in the gastrointestinal tract of humans and animals. Any food contaminated directly or indirectly with fecal material may theoretically contains one or more of these pathogens and thus can be potentially hazardous to consumers. They include *Escherichia coli*, *Enterococci* and *Clostridium perfringens* which are considered the most important fecal indicator pathogens in milk.

In recent years, *E. coli* has become recognized as a serious foodborne pathogen and has been associated with numerous outbreaks of disease in the UK, Japan and USA (Scotter *et al.*, 2000). *E. coli* includes a variety of different types that range from avirulent commensally strains that are present in the normal intestinal flora to highly virulent strains that cause a variety of sever infections in both humans and animals. Over 700 antigenic types or serotypes of *E. coli* had been recognized based on O, H and K antigens (Kaper *et al.*, 2004). Their pathogenicity

is considered to be mainly determined by specific virulence factors such as adhesions, invasions, toxins and capsule.

Raw milk and other dairy products, such as pasteurized milk and yoghurt contaminated with *E. coli*, have been the main cause of several outbreaks of milkborne diseases since the 1980s and thus remain a serious health risk (Seo *et al.*, 1998).

*Enterococcus*, formally the fecal or Lancefield group D *Streptococcus*, have been recognized to be of fecal origin since the beginning of this century and the usual ecological niche for *Enterococcus* species is the intestines of humans and animals. However, they are ubiquitous and can be found free living in soil, on plants, or in dairy products (Leclerc *et al.*, 1996). For many years, *Enterococcus* species were believed to be harmless to humans and considered unimportant medically because they produce bacteriocins and have been used widely over the last decade in the food industry as starter cultures (Foulquie Moreno *et al.*, 2006). In addition, they are present as active strains in different commercial probiotic preparations as a result of their ability to colonize the gastrointestinal tract and to control pathogenic bacteria (Lund *et al.*, 2002). However, recently *Enterococci* have been identified as secondary invaders in hospital-acquired infections (Murray, 1998) causing bacteraemia, endocarditis and other infections, with patient having a high mortality rate of up to 61% (Defátima Silva Lopes *et al.*, 2005). They are able to acquire genetic determinants for antibiotic resistances and among them the resistance to high levels of glycopeptides is of major concern (Bonten *et al.*, 2001). This dual nature of *Enterococci* gives rise to concern about the safety of using enterococcal probiotics or as starter cultures in the food industry (Lund and Edlund, 2001).

Different *Enterococcus* species are found in dairy products but *Ent. faecalis* and *Ent. faecium* remain the species of greatest importance (Gelsomino *et al.*, 2001). Moreover, *Enterococci* organisms have a distinctive role as an indicator of poor factory sanitation so that, they were enumerated and isolated from milk and milk products by many workers.

*Cl. perfringens* is widely distributed in nature occurring in soil, sewage, stool and intestinal tract of animals and humans (Steele and Wright, 2001). It is a classical agent of foodborne disease characterized by mildness and self limiting nature (Snaz *et al.*, 2002). The virulence of the organism is associated with the production of several toxins (exotoxin and enterotoxin). Moreover, the enterotoxigenic strains are a

common cause of food poisoning outbreaks worldwide (Labbe, 1991). *Cl. perfringens* enterotoxin (CPE), which is produced during the sporulation phase, causes the symptoms of *Cl. perfringens* food poisoning (Labbe, 1989). The illness is caused by ingestion of food containing a large number ( $>10^5$  cfu/g) of vegetative enterotoxigenic *Cl. perfringens* cells (Shandera *et al.*, 1983). The ingested bacteria multiply and sporulate, releasing CPE into the intestine causing watery diarrhea, abdominal pain, gas and necrotic enteritis in humans (Wise and Sirgusa, 2005).

*Cl. perfringens* has been isolated from dairy products and its presence could be attributed to contamination of raw milk used in production because its spores being heat resistant, and would not be destroyed during processing (Sinha and Sinha, 1986).

According to the public health hazard of fecal pollution microorganisms, there is a need to investigate milk and some dairy products sold in Assiut city for the incidence of *E. coli*, *Enterococci* and *Cl. perfringens*, so that the best indicator of direct fecal contamination or plant sanitation in milk and some dairy products could be developed.

## **MATERIALS and METHODS**

### **Samples:**

A total of 180 random samples of raw milk and some milk products including Kareish cheese, Domiati cheese, processed cheese, ice cream and yoghurt (30 samples each) were collected from various dairy shops, street vendors and supermarkets in Assiut city. All samples were collected in dry, clean and sterile containers and transferred to the laboratory with a minimum of delay. All samples were prepared for bacteriological examination according to APHA (1992).

### **Bacteriological examination:**

I- Coliforms, fecal coliforms and *E. coli*: Presumptive test for coliforms, confirmed test for coliforms, confirmed test for fecal coliforms and *E. coli* counts (MPN/ml or g) were carried out according to APHA (1992).

Identification of *E. coli* isolates: Morphological characteristic and biochemical tests were applied according to APHA (1992).

#### II- *Enterococci*:

- Enumeration of *Enterococci* was described by Deibel and Hartman (1982).

- Isolation and identification of *Enterococci* was carried out according to Morrison *et al.* (1997).

Identification of *Enterococci* isolates:

- Morphological characteristic: Microscopical examination (APHA, 1992) and motility test (Baron *et al.*, 1994).

- Biochemical tests: Catalase test (Finegold and Martin, 1982), growth at NaCl 6.5% (Sherman, 1937), sugar fermentation reaction (APHA, 1984), pyruvate fermentation (Gross *et al.*, 1975).

III- *Clostridium perfringens*:

Enumeration and identification of *Cl. perfringens* (MPN/ml or g) was done according to Beerens *et al.* (1980).

## RESULTS

**Table 1.** Incidence of fecal pollution indicators in the examined samples (no. =30).

Products	Coliforms		Fecal coliforms		<i>E. coli</i>		<i>Enterococci</i>		<i>Cl. perfringens</i>	
	Positive samples	%	Positive samples	%	Positive samples	%	Positive samples	%	Positive samples	%
Raw milk	30	100	18	60	1	3.3	25	83.3	-	-
Kareish cheese	30	100	25	83.3	10	33.3	25	83.3	1	3.3
Domiati cheese	7	23.3	2	6.7	1	3.3	8	26.7	-	-
Processed cheese	5	16.7	2	6.7	1	3.3	6	20	-	-
Ice cream	28	93.3	17	56.7	2	6.7	21	70	-	-
Yoghurt	14	46.7	11	36.7	6	20	12	40	-	-

**Table 2.** Frequency distribution of the positive samples based on their coliforms count.

Range	Raw milk		Kareish cheese		Domiati cheese		Processed cheese		Ice cream		Yoghurt	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
3-<10	-	-	1	3.3	3	42.8	2	40	2	7.2	5	35.7
10-<10 <sup>2</sup>	3	10	4	13.3	2	28.6	1	20	6	21.4	3	21.4
10 <sup>2</sup> -<10 <sup>3</sup>	-	-	2	6.7	1	14.3	1	20	6	21.4	1	7.2
10 <sup>3</sup> -<10 <sup>4</sup>	27	90	23	76.7	1	14.3	1	20	14	50	5	35.7

**Table 3.** Frequency distribution of the positive samples based on their fecal coliforms count.

Range	Raw milk		Kareish cheese		Domiaty cheese		Processed cheese		Ice cream		Yoghurt	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
3-<10	-	-	2	8	-	-	2	100	2	11.8	4	36.3
10-<10 <sup>2</sup>	3	16.7	2	8	2	100	-	-	6	35.3	1	9.1
10 <sup>2</sup> -<10 <sup>3</sup>	2	11.1	-	-	-	-	-	-	3	17.6	1	9.1
10 <sup>3</sup> -<10 <sup>4</sup>	13	72.2	21	84	-	-	-	-	6	35.3	5	45.5

**Table 4.** Frequency distribution of the positive samples based on their *E. coli* count.

Range	Raw milk		Kareish cheese		Domiaty cheese		Processed cheese		Ice cream		Yoghurt	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
3-<10	1	100	3	30	-	-	1	100	-	-	3	50
10-<10 <sup>2</sup>	-	-	-	-	1	100	-	-	1	50	1	16.7
10 <sup>2</sup> -<10 <sup>3</sup>	-	-	-	-	-	-	-	-	-	-	1	16.7
10 <sup>3</sup> -<10 <sup>4</sup>	-	-	7	70	-	-	-	-	1	50	1	16.7

**Table 5.** Frequency distribution of the positive samples based on their *Enterococci* count.

Range	Raw milk		Kareish cheese		Domiaty cheese		Processed cheese		Ice cream		Yoghurt	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
*<10 <sup>2</sup>	1	4	2	8	1	12.5	4	66.6	7	33.3	2	16.7
10 <sup>2</sup> -<10 <sup>3</sup>	5	20	1	4	2	25	-	-	1	4.9	3	25
10 <sup>3</sup> -<10 <sup>4</sup>	4	16	1	4	2	25	1	16.7	2	9.5	4	33.3
10 <sup>4</sup> -<10 <sup>5</sup>	8	32	2	8	1	12.5	1	16.7	2	9.5	1	8.3
10 <sup>5</sup> -<10 <sup>6</sup>	6	24	8	32	2	25	-	-	7	33.3	2	16.7
10 <sup>6</sup> -<10 <sup>7</sup>	1	4	11	44	-	-	-	-	2	9.5	-	-

\* No colonies could be detected on the plates but the organism could be isolated.

**Table 6.** Incidence of *Enterococci* strains recovered from the examined samples.

Isolated strains	Raw milk		Kareish cheese		Domiati cheese		Processed cheese		Ice cream		Yoghurt	
	No./30	%	No./30	%	No./30	%	No./30	%	No./30	%	No./30	%
<i>Ent. faecalis</i>	13	43.3	7	23.3	5	16.7	1	3.3	1	3.3	1	3.3
<i>Ent. faecium</i>	8	26.7	8	26.7	1	3.3	2	6.7	12	40	6	20
<i>Ent. hirae</i>	-	-	2	6.7	-	-	3	10	-	-	1	3.3
<i>Ent. durans</i>	-	-	3	10	-	-	-	-	1	3.3	-	-
<i>Ent. dispare</i>	1	3.3	1	3.3	-	-	-	-	1	3.3	-	-
<i>Ent. mundtii</i>	3	10	1	3.3	2	6.7	-	-	-	-	4	13.3
<i>Ent. saccharolyticus</i>	-	-	3	10	-	-	-	-	3	10	-	-
<i>Ent. cecorum</i>	-	-	-	-	-	-	-	-	2	6.7	-	-
<i>Ent. columbae</i>	-	-	-	-	-	-	-	-	1	3.3	-	-

## DISCUSSION

The obtained results summarized in Table 1 showed that coliforms, fecal coliforms, *E. coli* and *Enterococci* were detected in all the examined samples of milk and milk products. However, *Cl. perfringens* was recovered from 3.3% of the evaluated Kareish cheese samples only, while it failed to be detected in the other products.

The existence of coliforms and fecal coliforms in the examined raw milk samples were in parallel with those recorded by El-Zamkan (2007) and Abd El-Hameid (2002), respectively; while their presence in the examined Kareish cheese samples was in accordance to El-Zamkan (2007) and Ahmed *et al.* (1987), respectively.

From Table 2 it was clear that the majority of the examined Kareish cheese samples (96.7%) and ice cream samples (86.7%) did not comply with Egyptian Standards (2005a; 2005b) which recorded that coliforms must not exceed the level of 10 cfu/g. While, 90% of the examined processed cheese samples were in order with that standard. For Domiati cheese, 4 samples (13.3%) were above this standard. Egyptian Standards (2005c) reported that yoghurt must be free from coliforms, thus it was found that 46.7% of examined samples were not statutory with this standard and of inferior quality.

For fecal coliforms, Table 3 showed to what extent that raw milk, Kareish cheese, ice cream and yoghurt contained high frequencies in comparison with Domiati and processed cheeses.

Egyptian Standards (2005a, b, c and d) recorded that all the concerning samples must be free from *E. coli*. The obtained data

presented in Table 4 indicated that, 3.3% of raw milk, Domiati and processed cheese samples did not comply with these standards, while 33.3% of Kareish cheese samples also did not comply with it, fortunately, the majority 93.3 and 80% of ice cream and yoghurt samples respectively comply with its standards.

About *Enterococci*, It was clearly evident from Table 1 that *Enterococci* was existed in higher incidences in the concerning samples as compared with the level of *E. coli* or *Cl. perfringens*. This observation could be attributed to the greater resistance of *Enterococci* species to various unfavorable conditions such as salt content of Kareish and Domiati cheeses. As well as, the heat treatment applied during the manufacturing of processed cheese, freezing state of ice cream and low pH value of yoghurt. Consequently, greater resistance of *Enterococci* when compared with classical indicators of coliforms under unfavorable condition has led to an increasing tendency to include *Enterococci* in microbiological criteria as an indication of direct fecal contamination in various food products (Jay, 1992; Knudtson and Hartman, 1993; Audicana *et al.*, 1995).

In addition, there was an extremely low level of *Cl. perfringens* in contrast to higher rate of *E. coli* as demonstrated in the tested samples. This finding is in good agreement with the result estimated by Abd El-Rahman *et al.* (1994) that *E. coli* was able to mask the growth of toxigenic types of *Cl. perfringens*. This is because the ability of *E. coli* to produce colicins which have a role on the inhibition of other bacteria. Production of colicins from *E. coli* was also stated by other investigators (Djqnne, 1986; Ayhan and Aydin, 1989; Cong *et al.*, 1992).

Table 6 showed the incidence of the isolated *Enterococci* strains, in which, 9 strains were identified with high incidence in the examined raw milk, Kareish cheese and ice cream samples. *Ent. faecalis* and *Ent. faecium* were the most predominant strains. *Ent. faecalis* was isolated with high incidence 43.3% from raw milk followed by 23.3% from Kareish cheese whereas also *Ent. faecium* isolated with 40% from the examined ice cream samples.

In conclusion, the high level of *Enterococci* in the examined milk and milk products samples declared that *Enterococci* is preferable as fecal pollution indicator, while *E. coli* and *Cl. perfringens* should be included in food criteria as indicators of sanitation rather than fecal contamination. This result agreed to some extent with that recorded by Turantas (2002), Bahout and Moustafa (2006) who claimed that the levels of *Enterococci* were higher than fecal coliforms in ice cream and

frozen vegetables as well as Kareish cheese and thus favoring *Enterococci* as better indicators of contamination in such foods. Furthermore, this investigation indicated that the evaluated samples collected from the local markets in Assiut city were of poor sanitary quality which means fecal pollution and could pose a considerable risk to consumers.

## REFERENCES

- Abd El-Hameid, Karima G. (2002):* Studies on the sanitary conditions of raw milk in Qena Governorate. M.V.Sc. Thesis, Fac. Vet. Med., Assiut Univ., Egypt.
- Abd El-Rahman, M.; El-Bardisy, M.M.; Abd-El-Gaber, G. and Ahmed, H.F. (1994):* A study on *E. coli* and *Cl. perfringens* in buffaloe's milk. Giza. Vet. Med. J. 42(2):119-124.
- Ahmed, A.A-H.; Moustafa, M.K. and Abd El-Hakim, E.H. (1987):* Sanitary condition of Kareish cheese manufactured in Assiut city. Assiut Vet. Med. J., 19(37):76-80.
- APHA (American Public Health Association) (1984):* Compendium of Methods for the Microbiological Examination of Foods. 2<sup>nd</sup> Ed., Speck, H.L. (Ed.), Washington D.C., USA.
- APHA (American Public Health Association) (1992):* Compendium of Methods for the Microbiological Examination of Foods. 16<sup>th</sup> Ed., Washington D.C., USA.
- Audicana, A.; Perales, I. and Borrego, J.J. (1995):* Modification of Kanamycin-Esculin-Azid agar to improve selectivity in the enumeration of faecal *Streptococci* from water samples. Appl. Environ. Microbiol. 61:4178-4183.
- Ayhan, H. and Aydin, N. (1989):* Studies in the colicinogenic characteristics of *E. coli* strains isolated from man and animals. Etlik Veteriner Mikrobiyoloji Dergisi, 6(4):91-99.
- Bahout, A.A. and Moustafa, A.H. (2006):* Occurrence of some microorganisms in relation to public health in Kareish cheese. Assiut Vet. Med. J. 52(3):85-92.
- Baron, Ellen J.O.; Perteson, L.R. and Finegold, S.M. (1994):* Bailey and Scott's Diagnostic Microbiology. 9<sup>th</sup> Ed., Shanaton, J.F. (Ed.), Mosby-Year Book, Inc.
- Beerens, H.; Romond, Ch.; Lepage, C. and Criquelion, J. (1980):* A direct method for the enumeration of *Cl. perfringens* in foods

- and feces. World Congress Foodborne Infections and Intoxications, Berlin (West): 691-695.
- Bonten, M.J.; Willems, R. and Weinstein, R.A. (2001): Vancomycin-resistant *Enterococci*: why are they here, and where do they come from? *Lancet Infect. Dis.* 1:1314-1325.
- Cong, T.T.; Milon, A.; Boury, M. and Tasca, C. (1992): Colicins and resistance to colonization of *E. coli* 0103 enterocolitis of weaned rabbits. *Revue de Medecine Veterinaire*, 143(8/9):655-665.
- Defátima Silva Lopes, M.; Ribeiro, T.; Abrantes, M.; Figueiredo Marques, J.J.; Tenreiro, R. and Crespo, M.T.B. (2005): Antimicrobial resistance profiles of dairy and clinical isolates and type strains of *Enterococci*. *Int. J. Food Microbiol.* 103:191-198.
- Deibel, R.H. and Hartman, P.A. (1982): The *Enterococci*. In: *Compendium of Methods for the Microbiological Examination of Foods*. Speck, M.L. (Ed.), 2<sup>nd</sup> Ed., APHA. Inc.
- Djqnne, B.K., (1986): Colicin resistance in relation pathogenicity factors in strains of *E. coli* isolated from the intestinal tract of piglets. *Acta Vet. Scand.* 27(1):115-123.
- Egyptian Standards (2005a)*: Milk and Dairy products. Soft cheese. Egyptian Organization for Standardization and Quality Control. ES: 1008.
- Egyptian Standards (2005b)*: Milk and Dairy Products. Ice cream. Egyptian Organization for Standardization and Quality Control. ES: 1185.
- Egyptian Standards (2005c)*: Milk and Dairy Products. Yoghurt. Egyptian Organization for Standardization and Quality Control. ES: 1633.
- Egyptian Standards (2005d)*: Milk and Dairy products. Processed cheese. Egyptian Organization for Standardization and Quality Control. ES: 999.
- El-Zamkan, Mona A. (2007): Coliform bacteria in milk and some dairy products in Qena. M.V.Sc. Thesis, Fac. Vet. Med., South Valley Univ., Egypt.
- Finegold, S.M. and Martin, W.J. (1982): *Bailley and Scott's Diagnostic Microbiology*. 6<sup>th</sup> Ed., C.V. Mobsy Co., St. Louis, Toronto, London. UK.

- Foulquie Moreno, M.R.; Sarantinopoulous, P.; Tsakalidou, E. and De Vuyst, L. (2006):* The role and application of *Enterococci* in food and health. *Int. J. Food Microbiol.* 106:1-24.
- Gelsomino, R.; Vancanneyt, M.; ConDon, S.; Swing, J. and CoGan, T.M. (2001):* *Enterococci* diversity in the cheese-making environment of an Irish cheddar type cheese-making factory. *Int. J. Food Microbiol.* 71:177-188.
- Gross, K.C.; Houghton, M.P. and Senterfit, L.B. (1975):* Presumptive speciation of *Strep. bovis* and other group D *Streptococci* from human sources by using arginine and pyruvate tests. *J. Clin. Microbiol.* 1:54-60.
- ICMSF (International Committee on Microbiological Specification for Foods) (1978):* Microorganisms in Foods. I: Their significance and methods of enumeration. 2<sup>nd</sup> Ed., Univ. of Toronto, Press, Toronto Buffalo London.
- Jay, J.M. (1992):* Modern Food Microbiology, 4<sup>th</sup> Ed., pp.: 421-426. New York: AVI Books.
- Kaper, J.B.; Nataro, J.P. and Mobely, H.L.T. (2004):* Pathogenic *E. coli*. *Nat. Rev. Microbiol.* 2:123-140.
- Knudtson, L.M. and Hartman, P.A. (1993):* Comparison of fluorescent gentamicin-thallos-carbonate and KF Streptococcal agars to enumerate *Enterococci* and faecal *Streptococci* in meats. *Appl. Environ. Microbiol.* 59: 936-941.
- Labbe, R.G. (1989):* *Cl. perfringens*. In: Foodborne Bacterial Pathogens. M.P. Doyle (Ed.), Marcel Dekker, pp.: 191-234, New York, N.Y.
- Labbe, R.G. (1991):* *Cl. perfringens*. *J. Assoc. Off. Anal. Chem.* 74:711-714.
- Leclerc, H.; Devriese, L.A. and Mossel, D.A. (1996):* Taxonomical changes in intestinal (faecal) *Enterococci* and *Streptococci* consequences on their use as indicators of faecal contamination in drinking water. *J. Appl. Bacteriol.* 81:459-466.
- Lund, B. and Edlund, C. (2001):* Probiotic *Ent. faecium* strain is a possible recipient for the vanA gene cluster. *Clin. Infect. Dis.* 32:1384-1385.
- Lund, B.; Adamsson, I. and Edlund, C. (2002):* Gastrointestinal transit survival of an *Ent. faecium* probiotic strain administered with or without vancomycin. *Int. J. Food Microbiol.* 77:109-115.

- Morrison, D.; Woodford, N. and Cookson, B. (1997): *Enterococci* as emerging pathogens of humans. J. Appl. Microbiol., Symposium Supp. 83:89s-99s.
- Murray, B.E. (1998): Diversity among multidrug-resistant *Enterococci*. Emerg. Infect. Dis. 4:37-47.
- Oliver, S.P.; Jayarao, B.M. and Almeida, R.A. (2005): Foodborne pathogens in milk and the dairy farm environment: Food safety and public health implications. Foodborne Pathogens and Disease, 2 (2):115-129.
- Scotter, S.; Aldridge, M. and Capps, K. (2000): Validation of a method for the detection of *E. coli* O157: H7 in foods. Food Control, 11:85-95.
- Seo, K.H.; Brackett, R.E. and Frank, J.F. (1998): Rapid detection of *E. coli* O157:H7 using immuno-magnetic flow cytometry in ground beef, apple juice and milk. Int. J. Food Microbiol. 44:115-123.
- Shandera, W.X.; Tacket, C.O. and Blake, P.A. (1983): Food poisoning due to *Cl. perfringens* in the United States. J. Infect. Dis. 147:167-170.
- Sherman, J.M. (1937): The *Streptococci*. Bacteriol. Rev. 1:1-97 (Cited in ICMSF, 1978).
- Sinha, R.N. and Sinha, P.R. (1986): Incidence of *Clostridia* with reference to dairy processing. J. Food Sci. Technol. India, 25 (2):101-102.
- Snaz, J.C.; Dominguez, M.F.; Sagues, M.J.; Fernandez, M.; Feito, R.; Noguerales, R.; Asenio, A. and Fernandez Delahoz, K. (2002): Diagnosis and epidemiological investigation of an outbreak of *Cl. perfringens* food poisoning. Enferm. Inf. Microbiol. Clin. 20 (3):117-122.
- Steele, F.M. and Wright, K.H. (2001): Cooling rate effect on outgrowth of *Cl. perfringens* in cooked, ready-to-eat Turkey breast roasts. Poult. Sci. 6:813-816.
- Turantas, F. (2002): Incidence of faecal *Streptococci* as an indicator of sanitation in ice cream and frozen vegetables. Int. J. Food Sci. Technol. 37(3): 239-243.
- Wise, M.G. and Sirgusa, G.R. (2005): Quantitative detection of *Cl. perfringens* in the broiler foil gastrointestinal tract by real time PCR. Appl. Environ. Microbiol. 71:3911-3961.