

Dept. of Bacteriology,  
Animal Health Research Institute,  
Mansoura Provincial Laboratory.

## BACTERIOLOGICAL STUDY ON SOME CAUSES OF EARLY MORTALITY OF DUCKLINGS IN DAKAHLIA GOVERNORATE

(With 3 Tables)

By

**M. EL-SAYED HATAB; A.H. MOUSTAFA  
and M.M. ABD EL-LATIF**

(Received at 9/2/2005)

دراسة بكتريولوجية عن بعض مسببات النفوق المبكر في البط  
في محافظة الدقهلية

محمد السيد محمد حطاب ، عادل حسنين محمود مصطفى ،  
محمود محمد محمود عبد اللطيف

أجريت هذه الدراسة لمعرفة أسباب النفوق المبكر في البط. تم جمع ١٠٠ عينة من البط عمر (١-٢١ يوم) وأنواع مختلفة من عدة مزارع خاصة والعينات التي تصل إلى المعمل الفرعى بالدقهلية وكانت كلها تعاني من النفوق المبكر. تم فحص هذه العينات بكتريولوجيا وذلك بزرعها على أوساط مختلفة للبكتريا وقد أمكن عزل ١٤٤ عترة بكتيرية تم تصنيفها بالطرق المورفولوجية والبيوكيميائية إلى ٣٥ (٢٤,٣٠%) سيدوموناس أريجينوزا ، ٢٥ (١٧,٣٧%) إيشيريشيا كولاي، ٢٠ (١٣,٨٩%) لكل من السالمونيلا والإستافيلوكوكس أوريوس، ١٦ (١١,١١%) كليبيسيلا نيموني ، ١٥ (١٠,٤٢%) رايميريلأنا تيبتر و بروتيس فالجاريس ١٣ (٩,٠٢%). تم تصنيف معزولات الإيشيريشيا كولاي سيرولوجيا إلى ٧ عترات (O<sub>78</sub>K<sub>80</sub>(B-) و ٦ (O<sub>86</sub>K<sub>61</sub>(B7)) و ٥ (O<sub>125</sub>K<sub>70</sub>(B15)) و ٧ (untypable). كذلك تم تصنيف عترات السالمونيلا سيرولوجيا إلى ١٢ عترة سالمونيلا تيفيمورييم و ٨ عترات untypable. وقد أسفرت نتائج إختبارات حساسية المضادات الحيوية عن أن السيروفلووكساسين والإنروفلووكساسين والجنتاميسين كانت أكثر المضادات الحيوية تأثيراً على معظم المعزولات.

### SUMMARY

This study was conducted toward a problem of extend early mortality of ducklings. A total of 100 freshly dead and clinically sick ducklings of different ages (1-21 days) and types obtained from private farms at Dakahlia Governorate and also cases which arrive to the Mansoura lab.

The samples were dispatched to the laboratory to be examined bacteriologically for detection of the actual bacterial causes of early mortality problem in these farms. The obtained results pointed out that a total of 144 isolates were isolated. All the bacterial isolates were identified morphologically, culturally, biochemically and serologically for *E. coli* and *Salmonella* microorganisms. *Pseudomonas aeruginosa* was the most prevalent bacterial agent 35 (24.30%) followed by *E. coli* 25 (17.37) , *Salmonella* and *Staphylococcus aureus* 20 (13.89%) for each, *Klebsiella pneumoniae* 16 (11.11%), *Riemerella anatipestifer* 15 (10.42%) and *Proteus vulgaris* 13 (9.02%). The isolated *E. coli* were identified serologically into 7(O<sub>78</sub>K<sub>80</sub>(B-), 6(O<sub>86</sub>K<sub>61</sub>(B7), 5(O<sub>125</sub>K<sub>70</sub>(B15) and 7 (untypable), while the recovered *Salmonella* strains were identified into 12 *Salmonella typhimurium* and 8 (untypable). In-vitro-sensitivity pattern of isolated strains proved that ciprofloxacin, enrofloxacin, gentamycin were the most effective drugs.

*Key words: Mortality, ducklings.*

## INTRODUCTION

Nowadays, a great attention was payed toward ducks farming as a trial to fulfill excessive demand of the increased population from the animal protein. The ducks meat are considered to be of high protein content with high biological value.

Several microbial infections are responsible for the early mortality of ducklings and losses of duck industry. Many isolated microorganisms were associated with duck mortality as: *Salmonella*, *E.coli*, *Pseudomonas*, *Pasteurella*, *Proteus*, *Klebsiella* and *Staphylococci* (Bhowmik & Ray, 1987; EL-Gharib *et al.*, 1993 and Istanina, 1993). Antibiotics are used therapeutically and for prophylaxis in intensive domestic birds farming. However of bacteria resistant to antibiotics emerge even under control use of antibiotics (Helm *et al.* 1999). This study was done to investigate the possible bacterial causes of the problem.

## MATERIALS and METHODS

### A- Samples

A total of 100 freshly dead and clinical sick ducklings of different ages (1-21 days) and breeds were obtained from private farms and also cases which arrive Mansoura Provincial Laboratory. The



samples were dispatched to the laboratory without delay to be examined bacteriologically for isolation and identification of microorganisms agents.

#### **B- Clinical and PM examination:**

Ailing ducklings were examined clinically, then sacrificed and immersed in a disinfectant before being autopsied. Gross pathological changes were recorded, summarized and presented with results for both freshly dead & clinically sick ducklings.

#### **C- Media:**

- Nutrient agar media (Oxoid CM3).
- Blood agar media (Nutrient agar base Oxoid CM3 + 5-10% defibrinated sheep blood).
- Eosin methylene blue (Oxoid CM69).
- MacConkey agar (Oxoid CM7).
- Rappaport vassiliadis broth (Oxoid CM 669).
- Xylose lysine desoxycholate agar (Oxoid CM 469).

#### **D- Serological identification of Salmonella and E. coli:**

Samonella and E. coli isolates were identified according to (Edward and E wing, 1972).

#### **E- Bacteriological examination:**

Bacteriological samples were collected aseptically from the deep tissue of liver, lung, bone marrow, heart blood and brain. Each specimen was divided into 3 portions under aseptic condition. The first part was streaked onto predried surface of Blood agar, Nutrient agar, MacConkey agar (Oxoid, CM7) and Eosin methylene blue (EMB) (Oxoid, CM69), incubated aerobically at 37°C for 24 hours. The second part was inoculated into Rappaport Vassiliadis broth (RV) (Oxoid, CM 669) incubated at 42°C, after 24 hours incubation, loopfulls from R.V. enrichment were streaked onto Xylose lysine desoxycholate agar plate (XLD) (Oxoid, CM 469) with incubation at 37°C for 24 hours. The third part was streaked over blood agar plate containing 0.05% yeast extract, 5% newborn calf serum and 5 mg/100ml gentamycine, incubated in a candle Jar at 37°C for 24-48 hours.

The growing colonies on various plates were examined morphologically and biochemically as described by Baily & Scott, (1974); Cruickshank *et al.* (1975) and Carter (1984).

The identified E. coli strains were tested for enterotoxin production through grown the E. coli isolate in trypticase soya broth at 37°C in stationary culture overnight. Culture was centrifuged at 4000 rev/min. for 20 minutes. The supernatant was tested using commercially

VET-RPLA kits (reversed passive latex agglutination) from Oxoid (TD 0920 A) following the manufacturer's direction.

The biochemically identified *E. coli* and *Salmonella* isolates were subjected for serological identification using available *E. coli* test agglutinating sera (BioMerieux, 1986) and diagnostic *Salmonella* agglutinating antisera (Denka Selken Co. LTD, Tokyo, Japan) according to manufacturer's instruction.

**In vitro antibiotic sensitivity test:**

The disc diffusion technique was performed using Muller-Hinton medium (BioMerieux, France) on isolated bacteria from examined samples according to National Committee for Clinical Laboratory Standards (1984) and Quinn *et al* (1994).

## RESULTS

**The clinical signs and PM lesions:** The recorded clinical signs were diarrhea, ataxia, tremor of head and neck, coma, affected ducklings lie on their back, paddling their legs, septicemia. The P.M. examination of duckling revealed the presence of congestion in the internal organs (liver, spleen, intestine) and enlarged gall bladder.

**Bacteriological examination:**

The results of bacteriological examination were recorded in Tables 1, 2 and 3.

## DISCUSSION

Various clinical signs were recorded, diarrhea, ataxia, tremor of head & neck, coma, affected ducklings lie on their backs, paddling their legs, septicemia. Bacterial infection of the examined ducklings is one of the main causes of early mortality and the present study deals with the pathogenic bacteria responsible for early mortality of ducklings. *Pseudomonas aeruginosa*, *E.coli*, *salmonella*, *Riemerella anatipestifer*, *Kilebsiella pneumoniae*, *Proteus vulgaris* and *Staphylococcus aureus* were isolated in percentage of 24.3, 17.37, 13.89, 10.42, 11.11, 9.02 and 13.89 respectively. Similar findings were investigated by (El-Gharib *et al.* 1993 and Istanina, 1993).

*Pseudomonas aeruginosa* was the most prevalent bacterial agent 35 (24.3%). *Pseudomonas* infection of birds are of great important because epidemics may spread rapidly through flocks and is considered as one of the most dangerous disease, it causes morbidity and mortality



and the clinical signs including septicemia, diarrhea and respiratory signs (Soumaya, 1992 and Tanois & Samia, 1999).

Collibacillosis is a common systemic disease of world wide economic importance in birds, the major clinical signs of *E. coli* infection in young birds, acute septicemia that can cause sudden death (Leithnes and Heller, 1992).

In this study, all the isolated 25 (17.37%) *E. coli* strains from examined duckling samples were enterotoxigenic produce heat labile enterotoxin when tested by VET-RPLA kits, and serologically identified as 28% *E. coli* O<sub>78</sub> K<sub>80</sub> (B-), 24% *E. coli* O<sub>86</sub> K<sub>61</sub> (B7), 20% *E. coli* O<sub>125</sub> K<sub>70</sub> (B<sub>15</sub>) and 28% untypable strains (Table 2).

The results in Table (1) pointed out that *Salmonella* species were isolated from 20 (13.89%). This finding as nearly agree with the observations of El-Gharib *et al.* (1993). and Scott *et al.* (1984). The isolated *Salmonella* strains were serotyped as 12 (60%) *Salmonella typhimurium* and 8 (40%) untypable strains (Table2). *Salmonella typhimurium* had been reported as the most common species isolated from the examined ducks (50%) (Scott *et al.* 1984).

Generally, *Riemerella anatipestifer* is a Gram-negative, non-motile non sporforming rod, oxidase and catalase positive, not grow on MacConkey, negative indol and not hemolysis blood agar. *R. anatipestifer* is a contagious disease of domestic duck, it is known as duck septicemia, characterized by fibrinous pericarditis, perihepatitis, air sacculitis, caseous salpingitis and meningitis (Sandhu, 1986). Results in Table (1) showed that *Riemerella anatipestifer* were isolated in 15 (10.42%), our results nearly similar with Ziedler *et al.* (1984), reported high mortality in the fattening flocks of ducks occurred due to *Pasteurella anatipestifer* outbreaks. Regarding the isolation of *Klebsiella pneumoniae*, *Proteus vulgaris* and *Staphylococcus aureus*, they were isolated from the examined duckling samples at the incidence rate 11.11, 9.02 and 13.89% respectively. This findings similar to El-Gharib *et al.* (1993) and Istanina, (1993). The high mortality rate in duckling might be attributed to septicemic shock to the toxic effect associated with lipopolysaccharide fraction of the *Proteus* organisms (Wilson & Miles, 1975).

In vitro, the susceptibility distribution of each isolated pathogen to different antibiotics is presented in Table (3). The typical pattern of highly effective compounds were observed for zones of ciprofloxacin, enrofloxacin and gentamycin. These findings corresponded with those

reported by (Khodary & El-Sayed, 1997; Turbahn *et al.* 1997 and Rahman *et al.*, 1999).

In conclusion, the information given by the achieved results revealed that several microorganisms were incriminated in the early mortality of the ducks and therefore strict hygienic measurements should be applied on the egg laying, hatcheries, and good management during the production.

## REFERENCES

- Baily, W.R. and Scott, E.G. (1974): Diagnostic Microbiology. A Text Book for isolation and Identification of Pathogenic microorganisms. 4<sup>th</sup> Ed., the C.V. mosby Company Saint Louis.
- Bhowmik, M.K. and Ray M.M. (1987): Enteric diseases in ducks *Anas platyrhynchos Domesticus*. Ind. Vet. Med. J. 11(4): 209–214.
- BioMerieux (1986): Laboratory reagents and products bacteriology, Maclay, L' Etoile 69266, Charbonnieres, Les Bains, Frances.
- Carter, G.R. (1984): Procedures in Veterinary Bacteriology and Mycology. 4<sup>th</sup> Ed., Charles, G. Thomas, Puplicher Springfield Itlioms, USA.
- Cruickshank, R.; Duguid, J.P.; Morimon, B.P. and Swain, R.H.A. (1975): Medical Microbiology. 12<sup>th</sup> Ed., Vol. II, Churchill Livingstone. Edinburgth, London and New York.
- Edwards, P.R. and Ewing, MW.M. (1972): Identification of Enterobacteriaceae. 3<sup>rd</sup> Ed. Burgers Publishing Co. Minneoplos.
- El-Gharib, I.; Eldin – AMWK; Bastami, M.A.; Wahba, S.; Safwat, E.E.A. and Hatem, E. (1993): Incidence of isolation of microorganisms leading to embryonic mortalities and reducing hatchability of duck eggs. Vet. Med. J. – Giza, 41 (3): 63 – 65.
- Helm, J.D.; Hines, R.K.; Hill, J.E. and Caver, J.A. (1999): Multiple drug resistant *Salmonella typhimurium* DT 104 and DT 104 b isolated in Bobwhite quail (*colinus virginianus*). Avian Dis., 43: 788 – 791.
- Istania (1993): A bacteriological study on cases of post-hatching mortality in kedu chickens. Penyakit-Hewan. 25 (46): 124–127.
- Khodary, R.M. and El-Sayed, E.M. (1997): Treatment of ducklings salmonellosis by enrofloxacin. Assiut Vet. Med. J. 36: (72): 262–274.