

Dept. of Food Hygiene
Fac. Vet. Med., Alexandria University, Egypt

**PREVALENCE OF ENTEROHEMORRHAGIC
ESCHERICHIA COLI O157:H7 IN RAW MILK
AND EFFECT OF SOME CHEMICAL
PRESERVATIVES ON ITS VIABILITY**
(With One Table and 4 Figures)

By

A. A. AMER AND NABILA, F. SOLIMAN*

*Animal Health Research Institute, Alexandria,
Dept. of Food Hygiene
(Received at 8/5/2004)

مدى تواجد ميكروب الايشريكية القولونية المسببة للالتهاب المعوي النزلي
في اللبن الخام تأثير بعض المواد الحافظة على حيويتها

عمرو عبد المؤمن عامر ، نائلة سليمان

يعتبر اللبن غذاء صحي متكامل لكافة مراحل العمر وذلك لما يحتويه من عناصر غذائية هامة تجعله ذو قيمة عالية إلا أنه قد يتعرض للتلوث من مصادر مختلفة أثناء الإنتاج أو التداول مما يؤثر على جودته نتيجة تغيرات غير مرغوبة تجعله غير صالح للاستهلاك الأدمي. تم تجميع عدد ١٠٠ عينة من اللبن الخام من المزارع و محلات الألبان و الباعة الجائلين بمحافظة الإسكندرية والبحيرة لعزل ميكروب الايشريكية القولونية عترة O157:H7 المسببة للالتهاب المعوي النزلي. وقد اسفرت النتائج على تلوث العينات بميكروب الايشريكية القولونية عترة O157:H7 بنسبة ١%. وتم مناقشة الأهمية الصحية العامة للميكروب. كما تم دراسة تأثير بعض المواد الحافظة على حيويتها في اللبن المعقم مثل سوربات البوتاسيوم (٠,١٤٠,٠٥%) وبنزوات الصوديوم (٠,١٤٠,٠٥%) واستخدام سوربات البوتاسيوم وبنزوات الصوديوم معا بتركيز ٠,١% لكل منها في اللبن المعقم و نظام اللاكتوبيروكسينز المنشط في اللبن المعالج بالحرارة دون البسترة. ووجد ان بنزوات الصوديوم (٠,١%) لها القدرة على القضاء على الميكروب بعد ثمانية أيام من التخزين في درجة حرارة الثلاجة ووجد أن سوربات البوتاسيوم تقلل من العدد بمقدار لوغاريتم واحد فقط. ولكن عند استخدام سوربات البوتاسيوم (٠,١%) وبنزوات الصوديوم (٠,١%) معا فانها تقضي على الميكروب بعد ستة أيام من التخزين في درجة حرارة الثلاجة. أما بالنسبة الى نظام اللاكتوبيروكسينز المنشط في اللبن المعالج بالحرارة دون البسترة ووجد أنه يقلل من العدد بمقدار اثنان لوغاريتم.

SUMMARY

E. coli O157:H7, an Enterohemorrhagic pathogen, causes hemolytic uremic syndrome and hemorrhagic colitis in human. The objective of this study was determination of the prevalence of *E. coli* O157:H7 in raw milk and to evaluate the inhibitory effect of sodium benzoate, potassium sorbate, combination of sodium benzoate and potassium sorbate, and activated Lactoperoxidase system (LPS) on the viability of *E. coli* O157:H7 in commercially sterile milk stored at 4 °C. One hundred samples of raw milk were collected randomly from different areas in Alexandria and El-Behera Governorates, Egypt. The collected samples were examined for the prevalence of *E. coli* O157:H7 using Sorbitol MacConkey agar and rapid latex agglutination test for identification of *E. coli* serogroup O157. Different concentrations of Sodium benzoate (0.05 & 0.1%); Potassium sorbate (0.05 & 0.1%), combination of Sodium benzoate (0.1%) and Potassium sorbate (0.1%), and Lactoperoxidase system were added to detect their effects on *E. coli* O157:H7 viability. *E. coli* O157:H7 was detected in 1% of the examined samples. Sodium benzoate (0.1%) eliminates *E. coli* O157:H7 after 8 days and Potassium sorbate (0.1%) reduces *E. coli* O157:H7 one log cfu/ml after 10 days. The highest rates of inactivation occurred in the presence of the combination of Sodium benzoate and Potassium sorbate after 6 days, While LPS reduces *E. coli* O157:H7 2 log cfu/ml after 10 days. It seems necessary that the concerned authorities should impose regulation and bacteriological standards and take active part in the control of produced milk to ensure a maximum safety to the consumers.

Key words: *E. coli* O157:H7, milk, sodium benzoate, LPS, potassium sorbate.

INTRODUCTION

Milk is a highly perishable commodity and difficult to handle, especially in a country with high ambient summer temperature and lack of cooling methods.

E. coli O157:H7 constitutes a significance risk to human health world wide and the infection is associated with consumption of food of bovine origin (Philips *et al.* 2000), it causes acute renal failure in children (Fitzpatrick *et al.* 1991). The spectrum of clinical illness ranges from mild diarrhea, through bloody diarrhea and hemorrhagic uraemic

syndrome (HUS) and Thrombotic thrombocytopenic purpura (TTP) (Locking *et al.* 2001).

VTEC (verocytotoxin-producing *Escherichia coli*) O157:H7 has been identified as a possible contaminant of raw milk (Bryan, 1983). The gastro-intestinal tracts of ruminants, especially cattle, and humans are the main reservoirs of *E. coli* O157:H7 (Duffy *et al.*, (2001). Faecal contamination of raw cow's milk during its collection on dairy farms as well as milk filters are recognized as a major VTEC transmission route (Heuvelink *et al.*, 1998). Due to contamination of raw milk and/or improper processing, cow's milk has been implicated in foodborne outbreaks (Trevena *et al.*, 1996 and Keene *et al.*, 1997). In addition to foodborne transmission, *E. coli* O157:H7 may be spread by person to person as the source of infection is the feces of infected persons (principally sick persons, secondarily carriers) and objects contaminated by them (Griffin and Tauxe, 1991 and Dean-Nystrom *et al.*, 1997). *E. coli* O157:H7 constitutes 33% of milk borne general outbreaks of infectious intestinal diseases as a result of unpasteurized milk consumption (Gillespie *et al.*, 2003). Wachsmuth *et al.*, (1997) reported that raw milk was responsible for 5% of the outbreaks VTEC O157:H7 in the USA from 1982 to 1995.

Other verocytotoxin-producing serotypes (e.g. O26:H11, O103:H2, O113:H21) associated with human bloody diarrhea and HUS have also been isolated from sheep, calf and cattle faeces (Wells *et al.*, 1991).

Unhygienic conditions under which the animals are milked, small individual producer, long distance between the production and market areas, poor transportation, and insufficient or non-availability of milk cooling or chilling systems are the main problems of milk production. These problems lead to a considerable reduction in the shelf-life of milk consequently its souring (Jager, 1994 and Abd EL-Ghani and Sayed, 1997).

Preservatives are, generally, used to prevent or retard both chemical and biological deterioration of foods, to extend the shelf-life, retain nutritive value and to ensure the food safety by inhibiting or retarding the growth of pathogenic microorganisms. Addition of chemical preservatives to milk and milk products is widely used as an alternative method of cooling. Legal chemical preservatives are Sodium benzoate, Potassium sorbate and Lactoperoxidase system (Collins, 1971 and Ismail, 1997). *The objective* of this study was to determine the prevalence of *E. coli* O157:H7 in raw milk and to evaluate the inhibitory

effect of sodium benzoate, potassium sorbate, combination of sodium benzoate and potassium sorbate, and activated Lactoperoxidase system (LPS) on the viability of *E. coli* O157:H7 in commercially sterile milk stored at 4 °C

MATERIAL and METHODS

- I. **Prevalence of Enterohemorrhagic Escherichia coli O157:H7 in raw milk**
 1. **Collection and preparation of samples:**

One hundred samples of raw milk were collected randomly from different shops, street-vendors and supermarkets in Alexandria and El-Bchera Governorates, Egypt. Samples were transferred directly to the laboratory with a minimum of delay and were prepared according to APHA (1992).
 2. **Isolation and Identification of Escherichia coli O157:H7:**
 - 2.1. **Enrichment technique:** Milk samples were enriched in modified Tryptic Soya Broth (mTSB) supplemented with Novobiocin (20 mg/liter) and incubated at 37°C for 24 hours (DeBoor and Heuvelink, 2000).
 - 2.2. **Isolation on Sorbitol MacConkey agar:** Loopfuls from the incubated broth were streaked onto Sorbitol MacConkey agar plates and incubated at 37°C for 24 hours (DeBoor and Heuvelink, 2000).
 - 2.3. **Identification of Escherichia coli O157:H7:** Non Sorbitol fermenter colonies were identified biochemically as *E. coli* according to Varnam and Evans (1991) by the conventional IMViC, Urea hydrolysis and Triple sugar iron agar.
 - 2.4. **Identification of *E. coli* serogroup O157 (Oxoid,1996):** Typical *E. coli* strains were screened by a rapid latex agglutination test (*E. coli* O157, Oxoid diagnostic reagent 620 M) for identification of *E. coli* serogroup O157.
 - 2.5. **Identification of H7 (Difco, 1996):** Typical *E. coli* strains were screened by a tube agglutination test for identification of *E. coli* serogroup H7.
- II. **Effect of some chemical preservatives on the viability of Enterohemorrhagic Escherichia coli O157:H7 in milk**
 1. **Effect of different concentrations of Potassium sorbate and Sodium benzoate on viability of *E. coli* O157:H7 in commercial sterilized milk stored at refrigeration temperature at (4±1 °C):**

1.1. Organism: Culture of *E. coli* O157:H7 used in this study was obtained from the previously examined raw milk samples.

1.2. Experimental procedure:

1.2.1. Sterilized milk, which proved bacteriologically free from *E. coli* O157:H7, was inoculated with a suspension of 24 hours incubation *E. coli* O157:H7. The initial count was 1.8×10^3 cells/ml. Each inoculated milk sample was divided into 6 groups; Control (preservative free), A, B containing 0.05 and 0.1 % Potassium sorbate and C, D containing 0.05 and 0.1 % Sodium benzoate, respectively. While, group E contained a combination between Potassium Sorbate (0.1%) and Sodium benzoate (0.1%).

1.2.2. Effect of activated Lactoperoxidase system on viability of *E. coli* O157:H7 in thermalized milk stored at refrigerator temperature ($4 \pm 1^\circ\text{C}$):

Fresh milk samples were heated at 63°C for 10 minutes (Thermization) in flasks immersed in thermostatically controlled water bath. The flasks were inoculated with a suspension, of 24 hours incubation, *E. coli* O157:H7. The initial count was 1.8×10^3 cells/ml. The inoculated milk sample was divided into 2 parts; one was kept as a control while the second was activated with LP-system by adding 14 mg of Sodium Thiocyanate (Merck, Germany) followed by 30 mg of Sodium percarbonate (Aldrich, USA) /liter of milk (IDF, 1988).

Control samples together with inoculated ones were kept at $4 \pm 1^\circ\text{C}$ for studying the survival of *E. coli* O157:H7 on Sorbitol MacConkey agar (APHA, 1992). Counts were determined at Zero time, then examined every day till eleven days storage at refrigerator temperature ($4 \pm 1^\circ\text{C}$).

RESULTS

The obtained results are recorded in Table 1 and Figures 1-4.

DISCUSSION

I. Prevalence of Enterohemorrhagic *Escherichia coli* O157:H7 in raw milk

The result given in Table 1 revealed that out of 100 examined raw milk samples only one sample (1%) contained *E. coli* O157:H7.

This finding is in agreement with those reported by Steele *et al.*, (1997); Karakulska (2002); Murinda *et al.*, (2002) and Dontorou *et al.*, (2003). While higher incidences were reported by Clarke *et al.*, (1987)

and Awad (2002). On the other hand, Palmgren *et al.*, (1997), Aman *et al.*, (1998), Massa *et al.*, (1999) and Abd EL-Hady *et al.*, (2003) could not detect *E. coli* O157:H7 in any of examined raw milk samples. Reported estimates on the prevalence of VTEC O157:H7 in raw cow's milk ranged from 0 to 10% (Hancock *et al.*, 1998 and Coia *et al.*, 2001). However, according to surveys carried out in various countries, for example Scotland (Coia *et al.*, 2001) the Netherlands (Heuvelink *et al.*, 1998) and Sweden (Jonsson *et al.*, 2001), the prevalence of VTEC O157:H7 in raw cow's milk at farm level seems to be extremely low. Outbreaks of shiga toxin producing *E. coli* infection associated with serotypes O157:H7 or O157: H⁻ had been documented at least in 14 countries on 6 continents (C.D.C., 1993; Liesegang *et al.*, 2000 and Allerberg *et al.*, 2001).

Although over 150 different OH serotypes of VTEC have been associated with human illness, the main serotypes implicated in hemorrhagic colitis, HUS and thrombotic thrombocytopenic purpura are *Escherichia coli* O157:H7 and *E. coli* O157:H⁻ (Nelson *et al.*, 1998 and Vernozy-Rozand, 1999).

Nowadays verocytotoxin-producing *Escherichia coli* is considered an important emergent foodborne pathogen constituting a worldwide public health hazard either in the form of sporadic cases of infection or outbreaks (Vernozy-Rozand, 1999 and Leclercq *et al.*, 2001)

Moreover, the significant morbidity and mortality associated with VTEC infection largely as a result of hemolytic uremic syndrome (HUS) makes it a zoonotic problem of serious public health concern (Nelson *et al.*, 1998).

II. Effect of some chemical preservatives on the viability of Enterohemorrhagic *Escherichia coli* O157:H7 in milk.

a) Potassium sorbate

The results in Fig 1 reveal that a gradual decrease in numbers of *E. coli* O157:H7 occurred during the prolonged shelf-life of inoculated milk kept at 4±1°C. *E. coli* O157:H7 achieved a minimum count on the 7th and 10th day of storage before the samples spoiled at 8th and 11th day for samples containing 0.05 and 0.1% added Potassium sorbate, respectively. Potassium sorbate (0.1%) caused one log reduction in *E. coli* O157:H7 count by the end of the 10th day of storage. These findings agreed with those obtained by Narasimhan *et al.*, (1989) and El-Prince (1994) as they concluded that 0.1- 0.75% of Potassium sorbate inactivated *E. coli*. The inhibitory effect of Potassium sorbate on *E. coli* could be attributed to the reduction of intracellular pH of *E. coli* cells (Salmond *et al.*, 1984)

and the reduced uptake of required amino acids which are necessary for their growth (Ronning and Frank, 1987). Although the control samples were spoiled at 6th day as well as the treated samples, it is observed that Potassium sorbate greatly decrease the E. coli O157:H7 load in comparison with the control samples. E. coli O157:H7 survives or even multiplies in raw milk when stored at 8°C and reaffirm the need for pasteurization and holding the milk at $\leq 5^{\circ}\text{C}$ (Massa *et al.*, 1999 and Arias *et al.*, 2001).

b) Sodium benzoate

The data illustrated in Fig 2 verify that there was a steady decrease in numbers of E. coli O157:H7 during storage at refrigerator temperature ($4\pm 1^{\circ}\text{C}$) of inoculated milk samples containing 0.05% Sodium benzoate and reached a minimum count of 2.0×10^3 at 8th before the samples being spoiled. While in milk samples containing 0.1% Sodium benzoate, E. coli O157:H7 sharply decreases in numbers from 1.8×10^3 alls/ml to undetectable numbers by the end of the 8th day of storage.

c) Combination of Potassium sorbate (0.1%) and Sodium benzoate (0.1%)

It is evident from Fig 3 that E. coli O157:H7 was sharply decreased in inoculated milk during storage at ($4\pm 1^{\circ}\text{C}$). The combination could eliminate E.coli O157:H7 by the end of 6th day of storage. The obtained results verify that combination between Potassium sorbate (0.1%) and Sodium benzoate (0.1%) had a great inhibitory effect on E. coli O157:H7 in milk.

d) Effect of activated Lactoperoxidase system on viability of E. coli O157:H7 in thermalized milk.

The lactoperoxidase system (LPS) has been extensively investigated for dairy industry application (Farrag and Marth, 1992). When applied to dairy products, the major component of the system lactoperoxidase is present in the milk. The system is activated by the addition of thiocyanate and hydrogen peroxide in the form of sodium percarbonate. The activation of the LPS system in raw milk is used to prevent undue bacterial multiplication during collection and transport to the dairy processing plant in countries where refrigeration may not be feasible due to technical or economic reasons.

Fig 4 shows the inhibitory effect by the activation of LP-system on E. coli O157:H7 in milk stored at $4\pm 1^{\circ}\text{C}$. LP-system causes a 2 log reduction within 10 days before the samples were spoiled. Similar trends were observed by swart *et al.*, (1990) who reported that activation of LP-system in raw milk stored at 5°C for 4 days decreases E. coli count. Also,

Girgis *et al.*, (1999) mentioned that, the activation of LP-system in cow's milk caused a sharp decrease in the total coliforms including *E. coli* after 2 hours. The action of LP-system may be explained on the basis of production of bacterial inhibitor hypothiocyanate (OSCN) which interfere with metabolism of bacteria with regard to oxygen uptake and lactic acid metabolism thereby inhibiting growth (Wifson and summer, 1993 and De wit and Van Ilooydonk, 1996).

It was concluded from the results obtained that combination of Potassium sorbate (0.1%) and Sodium benzoate (0.1%) has more inhibitory effect on *E. coli*O157:H7 so, it is recommended that this combination should be added to raw milk to improve its shelf-life. Measures for improved cleanliness of stables, food management, efficient sanitation and cleanliness of animals when transported; the hygienic production of milk and milk products; strict maintenance of the cold chain (processing and distribution); heat treatment; provision of information to food handlers and to consumers with special attention to groups at special risk; and consideration of decontamination procedures before consumption should be applied.

REFERENCES

- A.P.H.A. American Public Health Association (1992):* Compendium of Methods for the Microbiological Examination of Foods, 2nd Ed. American Public Health Association, Washington, D.C., U.S.A.
- Abd El-Ghani, S. and Sayed, A. (1997):* Natural thiocyanate content and optimum condition for activation of Lactoperoxidase system in raw buffalo milk. *Egyptian J. Dairy Sci.*, 25:241.
- Abd El-Hady, H.M.; Essa, H.N.; and El-Essawy, H.A. (2003):* Rapid methods for detection of ENTEROPATHOGENIC AND ENTEROHEMORRHAGIC ESCHERICHIA COLI in some dairy products. The 2nd international congress of food Hygiene and human health 21 – 23 October, 2003. Fac. Vet. Med., Assiut Univ., Egypt.
- Allerberger, F.; Wagner, M.; Scheiger, P.; Rammer, HP.; Resch, A.; Dierich, MP.; Friedrich, AW. and Karch, H. (2001):* Escherichia coli O157:H7 and unpasteurized milk. *Euro. surveill.* 6(10):147-151.
- Aman, I.M., Knappstien, K. and Hahn, G. (1998):* Examination of Verotoxin-producing Escherichia coli in some Egyptian dairy products with special reference to serotype O157:H7. *Milchwissenschaft*, 53(12): 676-679.

- Arias, M.C.; Monge-Rojas, R.; Chaves, C. and Antillon, F. (2001):* Effect of storage temperature on growth and survival of *Escherichia coli* O157:H7 inoculated in foods from a tropical environment. *Rev. Biol.Trop.*49(2):517-523.
- Awad, E.I. (2002):* Studies on food poisoning bacteria with especial reference to campylobacter jejuni in milk. Ph. D. Thesis, Milk Hygiene. Fac. Vet. Med., Zagazig Univ., Egypt.
- Bryan, F., (1983):* Epidemiology of milk-borne diseases. *J. Food Prot.*, 46: 637-649.
- C.D.C. Center for Disease Control, (1993):* Update: multistate outbreak of *Escherichia coli* O157:H7. Infections from Hamburgers Western United States 1992-1993. *J. Am. Med. Assoc.* 269, 2194-2196.
- Clarke, R.C.; McEwen, S.A.; Gannon, V.P.; Valli, V.E.O.; Lior, H. and Gyles, C.L. (1987):* Isolation of Verotoxin-producing *Escherichia coli* from milk filters and calves in Ontario, Int. Symp. and Workshop on verocytotoxin producing infections. Abstract LFE.15.
- Coia, J., Johnston, Y., Steers, N., Hanson, M. (2001):* A survey of the prevalence of *Escherichia coli* O157:H7 in raw meats, raw cow's milk and raw-milk cheeses in south-east Scotland. *Int. J. Food Microbiol.*, 66: 63-69.
- Collins, E.B. (1971):* Preservatives in dairy food. *J. Dairy Science* 54:148-152.
- De Boor, E. and Heuvelink, A.E. (2000):* Methods for the detection and isolation of shiga toxin-producing *E. coli*. *J. App. Microbiol. Symposium Suppl.* 88,133-143.
- De Wit, J.N. and Hooydonk, A.C.M. (1996):* Structure, functions and application of lactoperoxidase in natural antimicrobial system. *Netherlands Milk Dairy J.* 50: 227.
- Dean-Nystrom, E.A.; Bosworth, B.T. and Moon, H.W. (1997):* Pathogenesis of O₁₅₇:H7 *Escherichia coli* infection in neonatal calves. *Adv. Exp. Med. Biol.* 412: 47-51.
- Difco (1996):* Serological identification of *Escherichia coli* O157:H7. Difco Lab. Detroit, Michigan, USA.
- Dontorou, C.; Papadopoulou, C.; Filloussis, G.; Economou, Y.; Apostolou, I.; Zakkas, G.; Salamoura, A.; Kansouzidou, A. and Levidiotou, S. (2003):* Isolation of *Escherichia coli* O157:H7 from foods in Greece. *Int. J. Food Microbiol.* 82(3): 273- 279.

- Duffy, G., Garvey, P., Wasteson, Y., Coia, J.E., McDowell, D.A. (2001): Epidemiology of Verocytotoxigenic *E. coli*. A technical booklet produced for an EU Concerted Action (CT98-3935). ISBN 1 84170 206 4.
- El-Prince, Enas, M. (1994): Effect of Potassium sorbate on the microbiological quality of milk and some milk products. Ph. D. Thesis. Milk Hygiene. Fac. Vet. Med., Assiut Univ., Egypt.
- Farrag, S. and Marth, E. (1992): *Escherichia coli* O157:H7, *Yersinia enterocolitica* and their Control in Milk by the Lactoperoxidase System: a Review. *Lebens-wiss. Und - Technol.* 25, 201-211.
- Fitzpatrick, M.; Shah, V. and Dillon, M. (1991): Long term outcome of childhood haemolytic uraemic syndrome. *Britain Med. J.* 303: 489-492.
- Gillespie, I.A.; G.K., Adak; O'Brien; S.J. and Bolton, F.T. (2003): Milk borne general outbreaks of infectious intestinal disease, England and Wales, 1992-2000. *Epidemiol. Infect.* 130(3):461-468.
- Girgis, E.; Abd El-Ghany, M.; Youssef, L. and Mohammed, L. (1999): Bactericidal effect of some pretreatment of raw milk on its keeping quality. *Egyptian J. dairy Sci.*, 27:59-70.
- Griffen, P.W. and Tauxe, R.V. (1991): The epidemiology of infection caused by *Escherichia coli* O157:H7, other enterohaemorrhagic *E. coli* and the associated hemolytic uremic syndrome. *Epidemiol. Rev.* 13, 60-98.
- Hancock, D.D., Besser T.E., Rice, D.H., Ebel, E.D., Herriott, D.E., Carpenter, L.V., (1998): Multiple sources of *Escherichia coli* O157 in feedlots and dairy farms in the Northwestern United States. *Prev. Vet. Med.*, 35: 245-250.
- Heuvelink, A., Bleumink, B., Biggelaar, F., Te Giffel, M., Beumer, R., De Boer, E., (1998): Occurrence and survival of verocytotoxin-producing *Escherichia coli* in raw cow's milk in the Netherlands. *J. Food Prot.*, 61: 1597-1601.
- International Dairy Federation, IDF (1988): Code of practice for the preservation of raw milk by the Lactoperoxidase system. Bull.No.234.
- Ismail, M.M. (1997): Some studies on milk and milk products preservation. M.V.Sc. Thesis, Fac. Agric., Mansoura Univ., Egypt.

- Jager, H. (1994): Effect of Nitrate on cheese ripening. *Milchwirtschaftliche Berichte aus den Bundesanstalten Wolfpassing und Rotholz*, 77:337-341.
- Jonsson, M., Aspan, A., Eriksson, E., Vagsholm, I. (2001): Persistence of verocytotoxin-producing *Escherichia coli* O157:H7 in calves kept indoors during the summer months in a Swedish dairy herd. *Int. J. Food Microbiol.*, 66: 55-61.
- Karakulska, J. (2002): Biochemical properties of enterotoxic and Enterohemorrhagic *E. coli* strain. *Med.DOSW. Mikrobiol.* 54(3): 215-223.
- Keene, W.E.; Hedberge, D.E.; Herriott, D.; Hancock, D.; Mackay, R.W.; Barrett, T.J. and Fleming, D.W. (1997): A prolonged outbreak of *Escherichia coli* O₁₅₇:H₇ infections caused by commercially distributed raw milk. *J. Infect. Dis.*, 176: 815-818.
- Leclercq, A.; Lambert, B.; Pierard, D. and Mahillan, J. (2001): Particular biochemical profiles for enterohaemorrhagic *Escherichia coli* O157:H7 isolates on the ID32E system. *J. Clin. Microbiol.* 39, 1161-1164.
- Liesegang, A.; Sachse, U.; Prager, R.; Claus, H.; Steinrück, H.; Aleksic, S.; Rabsch, W.; Voigt, W.; Fruth, A.; Karch, H.; Bockemühl, J. and Tschäpe, H. (2000): Colonal diversity of shiga toxin producing *Escherichia coli* O157:H7 / H⁻ in Germany – a ten year study. *Int. J. Med. Microbiol.* 290, 269-278.
- Locking, M.; Brien, S. and Reilly, E. (2001): Risk Factors for sporadic cases of *Escherichia coli* O157 infection : the importance of contact with animal excreta. *Epidemiol. Infect.* 127, 215-220.
- Massa, S.; Goffredo, C.; Allieri, C. and Natola, K. (1999): Fate of *Escherichia coli* O157:H7 in unpasteurized milk stored at 8 oC. *Let. Appl. Microbiol.* 28(1): 89-92.
- Murinda, S.E.; Nguyen, L.T.; Ivey, S.L.; Gillespie, B.E.; Almeida, R.A.; Draughon, F.A. and Oliver, S.P. (2002): Prevalence and molecular characteristic of *Escherichia coli* O157:H7 in bulk tank milk and fecal samples from cows; A 12- month survey of dairy farms in East Tennessee. *J. Food protection* 65(5); 725-729.
- Narasimhan, R.; Padmanaban, V.D. and Ulaganathan, V. (1989): Efficacy of some of the selected preservatives in sterile skim milk. *Indian Vet. J.* 66:51-55.
- Neaves, R., Deacon, J., Bell, C., (1994): A survey of the incidence of *E. coli* O157 in the UK dairy industry. *Int. Dairy J.*, 4: 679-696.

- Nelson, S.; Clarke, R.C. and Karmali, M.A. (1998): Verocytotoxin-producing *Escherichia coli* (VTEC) infection. In Palmer, S.R., Soulsby, L. and Simpson, D.I.H. Zoonoses. Biology, clinical practice and Public Health Control. Oxford University Press. pp. 89-104.
- Oxoid manual* (1996): Culture media, ingredient and other laboratory services. 15th Ed., Published by Oxoid Ltd., London.
- Palmgren, H.; Sellin, M.; Bergstorm, S. and Olsen, B. (1997): Enteropathogenic bacteria in migrating birds arriving in Sweden. *Scand. J. infectious. Diseases.* 29:565-568.
- Philips, A.; NavaBpour, S. and Hicks, S. (2000): Enterohaemorrhagic *Escherichia coli* O157:H7 target Peyer's patches in humans and cause attaching / effacing lesions in both human and bovine intestine. *Gut*, 47(3): 377-381.
- Rønning, I.E. and Frank, H.A. (1987): Growth inhibition of PA 3679 caused by stringent type response induced by protonophoric activity of sorbic acid. *Appl. Environ. Microbiol.* 53:1020-1027.
- Salmond, C.V.; Kroll, R.G. and Booth, I.R. (1984): The effect of food preservatives on PH homeostasis in *Escherichia coli*. *J. Gen. Microbiol.* 130: 2845-2850.
- Steele, M.L.; McNab, W.B.; Poppe, C.; Griffiths, M.W. and Chen, S. (1997): Survey on Ontario bulk tank raw milk for foodborne pathogens. *J. Food Prot.* 60(11):1341-1346.
- Swart, G., Jooste, P. and Mostert, J. (1990): The effect of the activated Lactoperoxidase system on raw milk bacteria. *Suid. Afrikaanse Tydskrif vir suiwelkunde* 22:29, *Dairy Sci., Abst.* 52:8126
- Trevena, W., Willshaw, G., Cheasty, T., Wray, C., Gallagher, J. (1996): Verocytotoxin-producing *E. coli* O157 infections associated with farms [letters; comment]. *Lancet*, 347: 60-61.
- Varnam, A.H. and Evans, M.G. (1991): Foodborne pathogens. An illustrated Textbook. Wolfe Publishing Ltd, New York. pp 129-155.
- Vernozy-Rozand, C. (1999): verotoxin-producing *Escherichia coli* (VTEC) and *Escherichia coli* O157:H7 in medicine and food industry. *Ann. Biol. Clin.* 57, 507-515.
- Wachsmuth, I.K., Sparling, P.H., Barrett, T.J., Potter, M.E., (1997): Enterohemorrhagic *Escherichia coli* in the United States. *FEMS, Immunol. Med. Microbiol.*, 18: 233-239.

Wells, J., Shipman, L., Greene, K., Sowers, E., Green, J., Cameron, D., Downes, F., Martin, M., Griffin, P., Ostroff, S., Potter, M., Tauxe, R., Wachsmuth, J., (1991): Isolation of *Escherichia coli* serotype O157:H7 and other Shiga-like-toxin-producing *E. coli* from dairy cattle. J. Clin. Microbiol., 29: 985-989.

Wilson, L.M. and Sumner, S.S. (1993): Antibacterial activity of the Lactoperoxidase system. A review J. Food Protection 65:887.

Table 1: Prevalence of *Escherichia coli* O157:H7 and *Escherichia coli* O157:H⁻ in raw milk samples.

	No. of samples	E. coli O157	E. coli O157:H7		E. coli O157:H ⁻	
			No.	%	No.	%
Raw milk	100	1	1	1	0	0




Figure 1: Effect of Potassium sorbate on viability of *E. coli* O157:H7 in sterile milk




Figure 2: Effect of Sodium benzoate on viability of *E. coli* O157:H7 in sterile milk




Figure 3: Effect of combination of Potassium sorbate (0.1%) and Sodium benzoate (0.1%) on viability of *E. coli* O157:H7




Figure 4: Effect of activated Lactoperoxidase system on viability of E. coli O157:H7 in thermalized milk.