

Dept. of Theriogenology
Fac. Vet. Med. Alex. Univ.

**EFFECT OF GnRH ON REPEAT BREEDER
BUFFALO-COWS AND DELAYED PUBERTAL
BUFFALO-HEIFERS AND ITS EFFECT WITH PGF₂ α
ON REPRODUCTIVE EFFECIENCY OF
POSTPARTUM BUFFALO-COWS**
(With 3 Tables)

By

**K.K. METWELLY; F.M. HUSSEIN; S.A. HATTAB
and F.K. KHALIFA**

(Received at 1/6/2004)

تأثير الهرمون المحرر للحاثة المنسلية على ظاهرة الشياح المتكرر
في الجاموس والبلوغ المتأخر في عجلات الجاموس ومصحوبا
بالبروستاجلاندين ف-2 ألفا على كفاءة التناسلية بعد الولادة

كمال متولى ، فكرى حسين ، سيد حطاب ، فوزى خليفة

تم إجراء ثلاثة تجارب الأولى استهدفت تأثير الهرمون المحرر للحاثة المنسلية GnRH عند اليوم 11 بعد التلقيح على ظاهرة الشياح المتكرر في الجاموس والثانية استهدفت دراسة تأثير GnRH على السيلوغ المتأخر في عجلات الجاموس أما الثالثة فقد استهدفت دراسة تأثير جرعتين من GnRH عند اليوم الأول والتاسع وجرعة من البروستاجلاندين ف-2 ألفا عند اليوم السابع وإجراء التلقيح عند اليوم العاشر (GPG) على خصوبة الجاموس بعد الولادة. فى التجربة الأولى تم استخدام 46 جاموسة ذات شياح متكرر تم حقن 34 منها بالهرمون المحرر للحاثة المنسلية عند اليوم 11 بعد التلقيح و تم حقن عدد 12 منها بمحلول ملح فسيولوجى. أما التجربة الثانية فقد تم استخدام عدد 41 من عجلات الجاموس أعمارها تعدت العامين ولم تظهر عليها علامات الشياح تم حقن عدد 30 منها بالهرمون المحرر للحاثة المنسلية وتم حقن 11 الباقيين بمحلول ملح فسيولوجى. وفى التجربة الثالثة تم استخدام عدد 50 جاموسة بعد الولادة بمسدة لا تقل عن 45 يوم تم حقن عدد 36 منها بجرعتين من الهرمون المحرر للحاثة المنسلية عند اليوم الأول والتاسع من التجربة وعند اليوم السابع تم حقنهم بالبروستاجلاندين ف-2 ألفا أما 14 جاموسة الباقيين فتم حقنهم بمحلول ملح فسيولوجى فى نفس المواعيد وتم تلقيح الخمسين جاموسة فى اليوم العاشر من التجربة. تم حساب عدد التلقيحات اللازمة للإخصاب والفترة من العلاج للإخصاب ومعدل الإخصاب فى التجارب الثلاث. أظهرت النتائج أن الهرمون المحرر للحاثة المنسلية أدى إلى تحسن الكفاءة التناسلية

للجاموس في التجارب الثلاث. ففي التجربة الثانية قد أظهرت النتائج أن عدد ٦، ٨٦ % من عجلات الجاموس المعالجة قد شاعت في حين أن ٥٤، ٥٠ % من المجموعة الضابطة قد شاعت. وفي المجموعة الأولى كانت عدد التلقيحات اللازمة للإخصاب في المجموعة المعالجة أقل منها في الضابطة. وكذلك أظهرت النتائج أنه في الثلاث تجارب كان هناك نقص معنوي في الفترة من العلاج للإخصاب وكذلك ارتفاع معنوي في معدل الإخصاب في المجموعات المعالجة مقارنة بالضابطة.

SUMMARY

Three experiments were conducted to study fertility of repeat breeder buffalo cows, delayed puberty in buffalo heifers and reproductive efficiency of postpartum buffalo cows after GnRH (at day 11 post service), GnRH and GnRH- PGF_{2α} - GnRII (GPG) administration in three separate experiments respectively. In experiment 1, 34 repeat breeder buffalo cows were injected (im) with 0.02 mg Buserelin (5 ml Receptal) at day11 post service (treated group). Other 12 repeat breeder buffalo cows were received (im) 5 ml saline and used as control group. In experiment 2, 30 delayed pubertal buffalo heifers (their ages above 2 years) were received (im) 0.02 mg Buserelin (treated group) however, 11 similar buffalo-heifers were injected (im) with 5 ml saline (control group). While in experiment 3, 36 of postpartum buffalo cows were received (im) 1st dose of GnRH (5ml Receptal) then injected (im) with PGF_{2α} at day 7 after the 1st dose of GnRH. At 40 hours post PGF_{2α} injection, animals received the second dose of GnRH (GPG treated group). Another 14 postpartum buffalo cows received saline and used as control. Cows in both GPG and control groups were naturally mated 24 hours after 2nd dose of GnRH or saline injections. Number of services per conception (S/C), treatment conception interval (TCI) and conception rate (CR) were calculated. The results indicated that, 26 (86.6%) out of 30 anoestrous buffalo heifers came in estrus after GnRH injection, while in control 6 (54.5%) out of 11 exhibited estrus signs. In addition, S/C was significantly lower ($P<0.05$) in treated than control animals in experiment 1. While, there were no significant differences in S/C between treated and none treated animals in experiments 2 and 3. There was a highly significant ($P<0.01$) reduction in TCI in treated (20.00±4.57, 21.68±3.06 and 23.62±3.26 days) than control (51.67±13.64, 46.40±8.06 and 49.13±7.62 days) in the three experiments respectively. Mean conception rate of the first 3 services was significantly higher (71, 80 and 72%) in the treated animals than in control (50, 66 and 57%) in the three experiments respectively. It was

concluded that, fertility parameters were improved after GnRH administration in repeat breeder buffalo cows, delayed puberty in buffalo heifers and in postpartum buffalo cows after GPG protocol.

Keywords: *GnRH, PGF_{2α}, repeat breeder, anoestrous, buffalo-cows*

INTRODUCTION

In buffalo-cows, poor reproductive performance is the major obstacle limiting their productivity and genetic progress. Although many variables can reduce fertility in buffaloes, the high incidence of silent ovulation (subestrus) either due to absence or reduced intensity of estrus behavior, or poor estrus detection is the most frequent cause of prolonged calving (Awasthi *et al.*, 1998; Barkawi *et al.*, 1998 and Singh *et al.*, 2000). One important approach to deal with reproductive inefficiency in buffaloes is the strategic use of hormonal agents to manage reproduction. Such programs are relatively recent in buffalo practice and their application is still and being refined in this species (Agarwal and Selvaraju, 2000 and Abou El-Roos and Abdel-Ghaffar, 2000).

GnRH is the key hormone that regulates the synthesis and release of LH and FSH from the anterior pituitary gland. The recent synthesis of GnRH has allowed its extensive use to initiate ovarian cyclicity (Roche and Mihm, 1996). The conception rate to first service after GnRH injection immediately following insemination was 53.4% compared with 32.33% in control non treated buffaloes (Rao and Rao, 1984). The cows that were injected with GnRH agonist on either day 11, 12 or 13 after insemination had higher pregnancy rates than control cows (72.4% versus 60.94%) (Macmillan *et al.*, 1986). GnRH treatment in cows on days 11 and 13 improved conception rate (Mann and Lamming, 1995). On the other hand, Jubb *et al.* (1990); Thatcher *et al.* (1993) and Ryan *et al.* (1994) were unable to show any improvement in conception rates when GnRH was injected between 11 and 13 days after insemination in repeat breeder cows.

Injection of anoestrous Egyptian buffaloes with GnRH improved their fertility (Nasr *et al.*, 1983, Hazzaa *et al.*, 1985, Aboul-ela *et al.*, 1988, Abdoon *et al.*, 1994 and El-Fadaly *et al.*, 1994). On the other hand, Khurana *et al.* (1982) found mild response after GnRH treatment in 3 of 21 anoestrous buffaloes. Similarly, Dhoble and Gupta (1988) reported that only 7 out of 34 anoestrous buffaloes expressed oestrus

after GnRH injection. Abeygunawardena *et al.* (1996) found that all treated anoestrous buffaloes with 2 GnRH injections 24 hours apart did not become pregnant within 90 days of treatment.

Combined use of GnRH and PGF₂ α in the postpartum period might be helpful for early resumption of ovarian activity in buffaloes, which have prolonged luteal activity (Shah *et al.*, 1990). Induction and synchronization of oestrus with acceptable level of fertility was obtained in anoestrous buffaloes receiving a GnRH followed by PGF₂ α (7 days interval) compared with untreated controls (Narasimha Rao and Venkatamiah, 1991). Conception rate was 77.8% after insemination in synchronized buffalo cows with PGF₂ α (Bicuda and Oba, 1992). Buffalo cows that were treated with GnRH- PGF₂ α responded in 80% ovulation without oestrus express that indicated by progesterone level above 1ng/ml (Hattab and Zeitoun, 1999). Administration of GnRH followed on day 7 by PGF₂ α elicited a rather poor response in induction of behavioral oestrus in anoestrous buffalo-cows (Narasimha Rao and Venkatamiah, 1991).

The present study aimed to investigate the effect of GnRH on reproductive performance of repeat breeder buffalo cows and delayed puberty in buffalo heifers and GnRH- PGF₂ α - GnRH (GPG) protocol on fertility of postpartum buffalo cows.

MATERIALS and METHODS

Animals:

The buffalo-cows and heifers that were used in the present study were selected sporadically from small holders in Etay El Barod district (El-Behira province). A total of 137 animals were used in this study. 96 buffalo cows, range in ages from 5 to 8 years and calved between October, 1999 and September, 2000. Other 41 buffalo heifers their ages were more than 2 years and they did not exhibit any signs of oestrus. The animals were kept in open shelters during daytime and were stabled by night all over the year. The feeding system during winter months included Trifolium plant (Barseem), rice straw and concentrate mixture. During summer months, Darawa was offered instead of Barseem. The animals were milked twice per day. Estrus signs were observed 3 times daily, at dawn, afternoon and evening for a period of 15 minutes per each time. The animals were naturally mated using fertile buffalo-bulls; pregnancy was diagnosed 60 days after mating by rectal palpation.

Treatment:

Experiment 1 (Repeat breeder):

In this experiment, 46 repeat breeder buffalo-cows were divided into two groups:

Treated group:

Thirty-four buffalo-cows were mated naturally during oestrus period. They were injected (im) with 5 ml Receptal (0.02 mg Buscrelin, a GnRH analogue manufactured by Intervet International B.V. Boxmeer – Holland) at day 11 post service.

Control group:

Twelve buffalo-cows received 5 ml saline at day 11 post natural mating.

Experiment 2 (anoestrous buffalo-heifers):

In this experiment, 41 delayed pubertal buffalo heifers, aged more than 2 years and their ovaries had no structures were divided into two groups.

Treated group:

Thirty delayed pubertal buffalo heifers were injected (im) with 5 ml Receptal.

Control group:

Eleven delayed pubertal buffalo heifers were injected (im) with 5 ml saline.

Fertile buffalo-bulls were used for natural mating of animals at oestrus.

Experiment 3 (GPG treated buffalo-cows):

This experiment was carried out on 50 lactating buffalo-cows, non pregnant during their postpartum periods (>45 days). The animals were free from any reproductive disorders. They divided into two groups.

Treated group:

Thirty six buffalo cows were treated according to the following Schedule (Jemmesonl, 1998).

At day 0: Animals were injected (im) with 1st dose of GnRH (5ml Receptal)

At day 7: Animals were injected by 2ml prosolvin (7.5 mg/ml Iuprostiol; a synthetic PGF₂ α analogue manufactured by Intervet International B.V. Boxmeer – Holland).

At day 9: Animals were injected (im) with 2nd dose of GnRH (5ml Receptal) 40 hours after PGF₂ α injection.

At day 10: All animals were naturally mated using fertile buffalo-bulls at fixed time 24 hours post 2nd dose of GnRH.

Control group:

Fourteen buffalo cows were given saline at the same time as those treated cows. These animals were naturally mated 24 hours post injection using fertile buffalo-bulls.

Number of services per conception (S/C), treatment conception interval (TCI) and conception rate (CR) were calculated in treated and control groups in each experiment.

Pregnancy diagnosis was carried out at day 60 post mating by rectal palpation.

Statistical analysis:

The data were analyzed by using SAS (1987). For analyzing the data of the treated and control group, T- independent test was used.

RESULTS

Results are presented in tables 1-3.

Experiment 1 (Repeat breeder buffalo-cows):

The effect of GnRH injection at day 11 post mating in repeat breeder buffalo-cows on S/C, TCI and CR are presented in table 1. The results indicated that, S/C was significantly ($P<0.05$) lower (1.61) in treated animal with GnRH than control (2.33). Moreover, GnRH injection resulted in a highly significant ($P<0.01$) reduction in TCI (20.00 ± 4.57 days vs. 51.67 ± 13.64 days) and significant increase ($P<0.05$) in the CR of the first three services (71% vs. 50%) in treated vs. control group respectively.

Experiment 2 (anoestrous buffalo- heifers):

Effect of GnRH on S/C, TCI and CR in anoestrous buffalo-heifers is presented in table 2. Twenty six (86.6%) out of 30 anoestrous heifers exhibited oestrus signs after treatment with GnRH. However, six (54.5%) out of 11 buffalo-heifers that injected by saline (control group) exhibited oestrus signs. S/C was non significantly lower (1.45) in treated animals than in control (1.8). The results also indicated that, there was a highly significant ($P<0.01$) reducing effect in TCI (21.68 ± 3.06 days) than control group (46.40 ± 8.06 days). In addition, CR of the first 3 services was significantly higher (80%) in treated animals than control group (60%).

Experiment 3 (GPG treated animals):

Results presented in table 3 indicate that, S/C in treated animals were not significantly lower (1.62) than in control animals (1.75). However, There was a highly significant ($P<0.01$) reduction in TCI in the GPG -protocol (23.62 ± 3.26 days) than control (49.13 ± 7.62 days).

Moreover, CR (72%) was significantly ($P < 0.05$) higher in treated animals than in control (57%).

Table 1: Effects of GnRH injection at 11 days post mating on S/C, TCI and CR in repeat breeder buffalo-cow (mean \pm SE)

Treatment	Number of buffaloes	S/C	TCI (days)	CR (%)			
				1 st service	2 nd service	3 rd service	Total
Treated	34	1.61	20.00 \pm 4.57 ^b	35.5	14.8	20.7	71 ^a
Control	12	2.33	51.67 \pm 13.64 ^a	8.4	25	16.6	50 ^b
T-value		1.893	2.791**				219.953*

Means in the same column have different superscripts are significantly different
 * $P < 0.05$, ** $P < 0.01$
 S/C = No. of services per conception
 TCI = Treatment conception interval
 CR = Conception rate

Table 2: Effects of GnRH injection on estrus exhibition, S/C, TCI and CR in anoestrous buffalo-heifers

Treatment	No. of buffalo heifers	Buffalo-heifers exhibited oestrus signs	S/C	TCI (days)	CR (%)			
					1 st service	2 nd service	3 rd service	Total
Treated	30	26 (86.6%)	1.45	21.68 \pm 3.06 ^b	53.2	19.2	7.6	80 ^a
Control	11	6 (54.5%)	1.80	46.40 \pm 8.06 ^a	33.0	16.5	16.5	66 ^b
T-value			0.862	3.865**				256.923*

Means in the same column have different superscripts are significantly different
 * $P < 0.05$, ** $P < 0.01$

Table 3: Effects of GPG protocol on S/C, TCI and CR in post partum buffalo-cows (mean \pm SE)

Treatment	Number of buffaloes	S/C	TCI (days)	CR (%)			
				1 st service	2 nd service	3 rd service	Total
Treated	36	1.62	23.62 \pm 3.26 ^b	38.8	22.2	11	72 ^a
Control	14	1.75	49.13 \pm 7.62 ^a	28.5	14.2	14.3	57 ^b
T-value		0.424	3.545**				225.203*

Means in the same column have different superscripts are significantly different
 * $P < 0.05$, ** $P < 0.01$

DISCUSSION

Reproductive inefficiency in buffalo-cows is a major problem in many herds and has been well documented (Gupta *et al.*, 1994; Dahama, 1995 and Singh *et al.*, 2000)

The reproductive parameters of the repeat breeder buffalo-cows that treated with GnRH at day 11 post service showed improvement where, S/C was reduced to 1.61 compared to control animals (2.33), TCI and CR were also improved in treated animals compared to control animals. These results are in agreement with that reported by Macmillan *et al.* (1986) and Mann and Lamming (1995). They reported that, administration of GnRH at day 11-13 after insemination in repeat breeder cows improved the pregnancy rates in treated animals. They suggested that GnRH treatment on days 11 and 13 causes a reduction in oestradiol secretion, which reduces the stimulus to the luteolytic mechanism, the chances of an embryo preventing luteolysis are increased and conception rate is improved.

On the contrary, Jubb *et al.* (1990); Thatcher *et al.* (1993) and Ryan *et al.* (1994) were unable to show any improvement in conception rates when GnRH was injected between 11 and 13 days after insemination in repeat breeder cows.

This study revealed that, GnRH injection in anoestrous buffalo-heifers resulted in 86.6% of buffalo-heifers came in oestrus, while in control, 54.5% of buffalo-heifers exhibited oestrus signs. An improvement was found in S/C, TCI and CR in treated compared with control. These results are in agreement with Pattabiraman *et al.* (1986). They found that, 60% of treated buffalo-heifers showed oestrus 10-26 days after treatment with GnRH while 20% from control animals came in oestrus by 35 days. Similarly, administration of GnRH in anoestrous Egyptian buffalo-cows improved their fertility (Nasr *et al.*, 1983, Hazzaa *et al.*, 1985, Aboul-ela *et al.*, 1988, Abdoon *et al.*, 1994 and El-Fadaly *et al.*, 1994).

On the other hand, Khurana *et al.* (1982) found mild response after GnRH treatment in 3 of 21 anoestrous buffalo-cows. Also, Dhoble and Gupta (1988) reported that, only 7 out of 34 anoestrous buffalo-cows expressed oestrus after GnRH injection. Abeygunawardena *et al.* (1996) found that, all treated anoestrous buffalo-cows with 2 GnRH injections 24 hours apart did not become pregnant within 90 days of treatment.

The close synchronization of ovulation produced by GPG technique reduced the need for oestrus detection and allowed optimal results with single fixed time insemination (Gordon *et al.*, 1996). The results in this study revealed that, there was a highly significant ($P < 0.01$) reduction in TCI and a significant ($P < 0.05$) higher CR in treated postpartum buffalo-cows with GPG protocol than in control animals.

These results are in agreement with that those published in cattle and buffaloes by Shah *et al.* (1990); Coleman *et al.* (1991); Guilbault *et al.* (1991) and Rao and Venkatramiah (1991). They reported that, a 6 or 7 days interval between GnRH and PGF₂α administrations was a satisfactory system for synchronization of oestrus with good fertility.

CR in GPG treated buffalo-cows in this study (72%) was nearly similar to that obtained by Bicuda and Oba (1992) who reported that, the conception rate was 77.8% after using GnRH immediately after insemination in synchronized buffalo-cows with PGF₂α. Moreover, Roche and Mihm (1996) reported that, GPG protocol induced synchronization of follicle waves with GnRH and control ovulation by using two days after PGF₂α.

El-Din Zain *et al.* (2001) and Metwelly and Abdelsabour-Elmaghraby (2001) concluded that, GnRH-PGF₂ α -GnRH is a useful synchronization method, as well as a tool for improving reproductive performance in Egyptian buffaloes.

On conclusion, GnRH administration at day 11 post service increased CR, improved S/C and reduced TCI in repeat breeder buffalo cows. It also improved fertility in delayed puberty of buffalo heifers. GPG protocol improved the fertility parameters of postpartum buffalo-cows.

REFERENCES

- Abdoon, A.S.S.; Ahmed, W.M.M. and El-Tohamy, M.M. (1994): Ovulation response following hormonal treatment of anoestrous buffaloes. In proceedings 4th world buffalo congress, San paulo, Barazil, 27-30 June, vol.3, 431-533.
- Abeygunawardena, H.; Kuruwita, V.Y. and Perera, B.M. (1996): Effect of exogenous hormones on fertility of postpartum anoestrous buffaloes. In rule of the buffaloes in rural development in Asia. Proceedings of a regional symposium, Peradeniya Siri Lanka, 10-15 December 1995 (edited by Perera, B.M.A.O.; Siriwardene, J.A. de S., Horadagoda, N.U. and Ibrahim, M.N.M.) Colombo 7, Siri Lanka; ARESA Press, 1996: 327-349.
- Aboul-ela, M.B.; Khattab, R.M.; El-Keraby, F.E.; Shafie, M.M. and Bedier, L.H. (1988): Patterns of ovarian and oestrus activity and induction of cyclic activity during the postpartum period in Egyptian buffaloes. Proceedings of the final research

- coordination meeting on an optimizing grazing animal productivity in the Mediterranean and North African regions with aid of nuclear techniques. March 23-27, Rabat, Morocco, PP. 239-253.
- Abou El-Roos, M.E.A. and Abdel Ghaffar, A.E. (2000):* Some trials to improve the reproductive efficiency of subestrus buffalo cows using PGF 2α at midluteal phase. *Assiut Vet. Med. J.*, 43(86): 327-337.
- Agarwal, S.K. and Selvaraju, S. (2000):* Recent technologies to improve reproductive efficiency of cattle and buffaloes. *Advancement in Veterinary Sci., Ind. Vet. Cong. Izatnyar, India*, 18-19 Feb. P. 96.
- Awasthi, M.K.; Tiwari, R.P. and Pangaonkar, G.R. (1998):* Induction of estrus and fertility with low dose of prostaglandin F 2 alpha in subestrus buffaloes. *Ind. J. Anim. Sci.*, 68 (10): 1049-1050.
- Barkawi, A.H.; Khattab, R.M. and El-Wardani, M.A. (1998):* Reproductive efficiency of Egyptian buffaloes in relation to estrous detection systems. *Anim. Reprod. Sci.*, 51 (3): 225-231.
- Bicuda, S.D. and Oba, E. (1992):* The use of gonadotrophin in oestrus synchronization programs in buffalo for artificial insemination: Preliminary investigation. *Ind. J. Dairy Sci.*, 45 (1): 1-2.
- Coleman, D.A.; Bartol, F.F.; Spencer, T.E.; Floyd, J.G.; Wolfe, D.F. and Brendemuhl, J.P. (1991):* Effects of a potent GnRH agonist on hormonal profiles, synchronization of oestrous and fertility in beef cattle. *J. Anim. Sci.*, (suppl.1): 396 (Abst).
- Dahama, R.S. (1995):* Genetic analysis of reproductive traits in buffaloes. *Indian J. Dairy Sci.*, 48 (5): 317-322.
- Dhoble, R.L. and Gupta, S.K. (1988):* Response to synthetic gonadoliberin (GnRH) in anoestrous buffaloes. *Ind. J. Anim. Sci.*, 58 (9): 1071-1072.
- El-Din Zain, A.; Abdel-Razek, A.Kh. and Anwar, M.M. (2001):* Effect of combined using of GnRH and PGF 2α on estrus synchronization and pregnancy rate in buffalo cows. *Assiut Vet. Med. J.*, 45(89): 303-316.
- El-Fadaly, M.A.; Atiefa, A.A.; Abass, H.I.; El-Essawy, G.S. and El-Essawy, S.A. (1994):* Induction of cyclicity in anoestrus and subestrus Egyptian buffaloes. *Proceedings, 4th world buffalo congress, Sao Paulo, Barazil, 27-30 June, 1994: Volume3, 549-551.*

- Gordon, P.G.; Peters, A.R.; Ward, S.J. and Warren, M.J. (1996):* The use of prostaglandin in combination with a GnRH agonist in controlling the timing of ovulation in dairy cows. BCVA Edinburgh, Vol.1, 164-169.
- Guilbault, L.A.; Villeneuve, P.; Laverdiere, G.; Proulx, J. and Dufour, J.J. (1991):* Oestrous synchronization in beef cattle using a potent GnRH analogue (buserelin) and cloprostenol. *J. Anim. Sci.*, 69 (Suppl.1): 419.
- Gupta, B.D.; Kaushik, S.N. and Mishra, R.R. (1994):* Study on reproduction efficiency parameters of Murrah buffaloes. *Indian J. Dairy Sci.*, 47 (4): 257-264.
- Hattab, S.A. and Zeitoun, M.M. (1999):* Progesterone profile response to exogenous FSH, PMSG or GRH-PGF₂ α in non cycling buffalo cows and effect of FSH or PMSG on early pregnancy. *Alex. J. Agric.*, 44 (3): 21-40.
- Hazzaa, A.M.; Abdel-Raheim, A. and El-Ghannam, F. (1985):* Trials on the treatment of inactive ovaries in cows and buffaloes with GnRH (Hoe, 7666). *Zag. Vet. Med. J.*, 33 (1) 297-303.
- Jubb, T.F.; Abhayaratne, D.; Malmo, J. and Anderson, G.A. (1990):* Failure of an intramuscular injection of analogue of gonadotrophin releasing hormone (Buserelin) at 11-13 days post insemination to increase pregnancy rates in dairy cattle. *Aust. Vet. J.*, 67: 359-361.
- Khurana, K.; Tyagi, R.P.S.; Gupta, R.C. and Verma, S.K. (1982):* Gonadotrophin releasing hormone in the treatment of anoestrous buffalo cows and heifers. *Ind. Vet. J.*, 59: 479-480.
- Macmillan, K.L.; Taufa, V.K. and Day, A.M. (1986):* Effects of an agonist of gonadotrophin releasing hormone (Buserelin) in cattle. III pregnancy rates after a post insemination injection during metestrous or dioestrous. *Anim. Reprod. Sci.*, 11: 1-10.
- Mann, G.E.; and Lamming, G.E. (1995):* Effects of treatment with buserelin on plasma concentrations of oestradiol and progesterone and cycle length in the cow. *Br. Vet. J.*, 151, 427-429.
- Metwally, K.K. and Abdelsabour-Elmaghraby, M.M. (2001):* Reproductive performance of lactating Egyptian buffaloes treated with Gonadotropin-Releasing hormone and/or Prostaglandin F₂ α for ovulation or estrus synchronization. 6th Sci. Cong. Egyptian Society for Cattle Diseases, 4-6 Nov. 2001. Assiut, Egypt (250-258).

- Narasimah Rao, A.V. and Venkatramiah, P. (1991):* Induction and synchronization of Oestrous and fertility in seasonally anoestrous buffaloes with GnRH and PGF analogue. *Anim. Rep. Sci.*, 25: 109-113.
- Nasr, M.T.; Sharawy, S.; El-Azab, M.A. and Labib, F.M. (1983):* Induction of oestrus and improvement of fertility in anoestrous cow and buffaloes with Receptal. *The blue book* 32: 91-93.
- Pattabiraman, S.R.; Veerapandian, C. and Quayam, S.A. (1986):* Effects of Receptal treatment in anoestrous and early postpartum cows and buffaloes. *Indian Vet. J.*, 63: 409-413.
- Rao, A.R. ad Rao, K.S. (1984):* Improved conception rate in buffaloes after administration of Receptal (Buserelin). *Ind. Vet. J.*, 61(9): 8-13.
- Rao, A.V.N. and Venkatramiah, P. (1991):* Induction and synchronization of estrus and fertility in seasonally anoestrous buffaloes with GnRH and a PGF_{2α} analogue. *Anim. Rep. Sci.*, 25: 109-113.
- Roch, J.F. and Mihm, M. (1996):* Physiology and practice of induction and control of oestrus in cattle. *BCVA Edinburgh* 157-183.
- Ryan, D.P.; Snijders, S.; Condon, T.; Grealy, M.; Sreenan, J. and O'Farrell, K.J. (1994):* Endocrine and ovarian responses and pregnancy rates in dairy cows following the administration of gonadotrophin releasing hormone analogue at the time of artificial insemination or at mid-cycle post insemination. *Anim. Reprod. Sci.*, 34: 179-191.
- SAS (1987):* Statistical Analysis System. User^o guide: Statistics., SAS institute, Cary, North Carolina.
- Shah, N.S.; Willemsse, A.H. and Van de Wiel, D.F.M. (1990):* Reproductive performance of Nili-Ravi buffaloes after a single injection of GnRH early postpartum. *Trop. Anim. Hlth. Prod.*, 22: 239-246.
- Singh, J.; Nanda, A.S. and Adams, G.P. (2000):* The reproductive pattern and efficiency of female buffaloes. *Anim. Reprod. Sci.*, 60-61: 593-604.
- Thatcher, W.W.; Drost, M.; Savio, J. D.; Macmillan, K.L.; Entwistle, K. W.; Schmitt, E.J.; Dela Sota, R.L. and Morris, G.R. (1993):* New clinical uses of GnRH and its analogue in cattle. *Anim. Reprod. Sci.*, 33: 27-49.