

Animal Health Research Institute,
Alexandria Branch

PREVALENCE OF PSYCHROTROPIC FOOD BORNE PATHOGENS IN FISH IN ALEXANDRIA MARKETS

(With 4 Tables)

By

AMAL.F.A. MANSOUR and F.A. EL-SHABOURY

(Received at 16/3/2009)

مدي تواجد الميكروبات الممرضة المحبة للبرودة في الأسماك في أسواق
محافظة الإسكندرية

آمال فهمي علي منصور ، فتح الله علي الشابوري

تم جمع 100 عينة من الأسماك الطازجة عبارة عن 50 عينة من أسماك البلطي و 50 عينة من أسماك القرموط وذلك لتقييمها من الناحية الصحية وبإجراء الطرق القياسية لتقدير العد البكتيري للميكروبات المحبة للبرودة ففي أسماك البلطي دلت النتائج علي أن متوسط العدد الكلي للميكروبات المحبة للبرودة $3.2 \times 10^2 \pm 1.3 \times 10^2$ /جرام بينما كان متوسط العدد الكلي لميكروب السودوموناس $4.5 \times 10^3 \pm 1.3 \times 10^3$ /جرام ومتوسط العدد الكلي لميكروب الباسيلس سيرس $3 \times 10^3 \pm 3.6 \times 10^2$ /جرام وكانت أعلى نسبة للتلوث في اسماك البلطي تحتوي علي الميكروبات المحبة للبرودة تتراوح بين 10^4 و 10^5 بينما أعلى نسبة تلوث بميكروب السودوموناس والباسيلس سيرس تتراوح بين 10^3 و 10^4 . وفي أسماك القرموط دلت النتائج علي أن متوسط العدد الكلي للميكروبات المحبة للبرودة $2.5 \times 10^4 \pm 3.6$ × 10^3 /جرام ومتوسط العدد الكلي لميكروب السودوموناس $6.8 \times 10^3 \pm 1.2 \times 10^3$ /جرام ومتوسط العدد الكلي لميكروب الباسيلس سيرس $8.6 \times 10^2 \pm 2.1 \times 10^2$ /جرام. وكانت أعلى نسبة تلوث في الأسماك المحبة للبرودة تتراوح بين 10^4 و 10^5 بينما أعلى نسبة للتلوث بميكروب السودوموناس تتراوح بين 10^3 و 10^4 وأخيراً التلوث بميكروب الباسيلس سيرس يتراوح بين 10^2 و 10^3 .

SUMMARY

One hundred samples of fish represented by 50 samples of *Tilapia nilotica* and 50 samples of *Claris lazera* were collected from different fish markets in Alexandria Governorate to evaluate their microbiological quality and detection of food borne psychrotrophs. The bacteriological examination revealed that the mean values of total psychrotrophic count, total *Pseudomonas* count and total *Bacillus cereus* count in *Tilapia nilotica* were $3.2 \times 10^2 \pm 1.3 \times 10^2$, $4.5 \times 10^3 \pm 1.3 \times 10^3$ and $3 \times 10^3 \pm$

3.6×10^2 respectively. The majority of *Tilapia nilotica* fish samples had psychrotrophic count ranged from 10^4 to $<10^5$. While, the majority of such examined fish samples had *Pseudomonas* and *Bacillus cereus* count between 10^3 to 10^4 . Furthermore, the average counts of psychrotrophics, *Pseudomonas* and *Bacillus cereus* for *Claris lazera* were $2.5 \times 10^4 \pm 3.6 \times 10^3$, $6.8 \times 10^3 \pm 1.2 \times 10^3$ and $8.6 \times 10^2 \pm 2.1 \times 10^2$ respectively. The public health significance of the isolated microorganisms and possible sources of contamination of fish with such organisms as well as some recommendations to improve the quality of fish were discussed.

Key words: *Fish, Tilapia nilotica, Claris lazera, psychrotrophs, B-cereus, Pseudomonas spp.*

INTRODUCTION

Fish and fish products were incriminated as causes of food poisoning, food intoxication, allergic and skin disorders as well as other many infectious diseases (Lawsan, 1970). Microorganisms, individually and as a group, grow over a very wide range of temperatures. Therefore, it would be well to consider at this point the temperature growth ranges for organisms of importance in foods as an aid in selecting the proper temperature for the storage of different types of foods.

Psychrotrophs are those organisms that grow well at or below 7°C and have their optimum between 20°C and 30°C . *Bacillus cereus* and species of *Pseudomonas* were found among the psychrotrophic microorganisms, and their presence or contamination of food creates a great risk as they lead to food poisoning and/or spoilage of food products (Jay, 2000). In addition (Parry *et al.*, 1983) mentioned that *Bacillus cereus* causes health hazard to consumers, and also important for economic as they cause spoilage and deterioration of meat and meat products and responsible for food poisoning outbreaks among the consumers. Refrigeration is often the main and frequently the only factor to control food-borne pathogens in these types of foods. Hence, temperature abuse of such foods can result in food-borne illness. In addition, some psychrotrophic pathogens can grow in refrigerated foods with little or no obvious change of sensory characteristics (Berrang *et al.*, 1989). *Bacillus cereus* can give rise to two distinct forms of food-borne disease, the emetic forms and the diarrhaeal syndromes. The emetic form believed to be associated with an emetic toxin performed in food while the diarrhaeal type is caused by an enterotoxin (Altayer and Sutherland, 2006).

Pseudomonas species were responsible for spoilage and the identified off-odours, odours produced by the spoilage *Pseudomonas* bacteria were varied that resemble sulphur, ammonia and bad odour (Russel *et al.*, 1995) *Pseudomonas* microorganisms were isolated and determined as causes for contamination and spoilage of fishes (Gram and Melchiorson, 1996; Yumoto *et al.* 1999; Carraseal *et al.* 2002). The wide spread occurrence and psychrotrophic nature of these bacteria could increase the risk as refrigerated ready-to-eat foods may serve as vehicles of food-borne illness. The aim of this study was to assess the presence of some psychrotrophic microorganisms in ready-to-eat fishes. The health risks due to these pathogens for consumers were also discussed.

MATERIALS and METHODS

Collection of samples:

A total of 100 random ready-to-eat fish samples, 50 each of *Tilapia nilotica* and *Claris lazira* were collected from fish markets in Alexandria city. The collected samples were transferred in an ice box immediately to the laboratory under complete aseptic precaution without undue delay for bacteriological examination.

Preparation of samples (APHA, 1984)

10 grams of each of fish and 90 ml of sterile peptone water were added and thoroughly mixed by using sterile blender for approximately 2.5 minutes, from which ten fold serial dilutions were prepared. The prepared samples were subjected to the following examination.

1- Determination of psychrotrophic bacterial count:

Standard plate count agar was used as recommended by APHA (1992). The average number of colonies per gram was determined and the psychrotrophic count/g samples was calculated and recorded.

2- Enumeration and identification of *Pseudomonas* species (ICMSF, 1978):

0.1 ml. of each fish muscle dilutions was separately inoculated into duplicated Petri-dishes of *Pseudomonas* selective agar medium supplemented with glycerol. The inoculated plates were incubated at 25°C for 48 hours after which all developed colonies (greenish yellow pigment) were enumerated and the average count/g was calculated and recorded.

3-Enumeration and identification of *Bacillus cereus* (ISO, 1987):

The spreading technique was applied on the surface of *Bacillus cereus* selective agar which incubated at 37°C for 24 hours, then the count was recorded. Colonies though to be *Bacillus cereus* were identified by microscopical examination and biochemical reactions.

RESULTS

Table 1: Statistical analysis of total psychrotrophic, *Pseudomonas* and *Bacillus cereus* of the examined *Tilapia nilotica* (n=50).

Bacterial count	Total Psychrotrophic	<i>Pseudomonas</i>	<i>Bacillus cereus</i>
Minimum	1.7X10 ³	7X10 ²	7.8X10 ²
Maximum	3X10 ⁴	2.6X10 ⁴	8X10 ³
Mean	3.2X10 ²	4.5X10 ³	3X10 ³
Standard error	1.3X10 ²	1.3X10 ³	3.6X10 ²

Table 2: Frequency distribution of the examined *Tilapia nilotica* based on their total psychrotrophic, *Pseudomonas* and *Bacillus cereus* counts (n=50).

Count Range	Total Psychrotrophic		<i>Pseudomonas</i>		<i>Bacillus cereus</i>	
	No.	%	No.	%	No.	%
< 10 ²	25	50	27	54	28	56
10 ² < 10 ³	0	0	3	6	4	8
10 ³ < 10 ⁴	10	20	12	24	18	36
10 ⁴ < 10 ⁵	15	30	8	16	0	0

< 10² mean negative result.

Table 3: Statistical analysis of total psychrotrophic, *Pseudomonas* and *Bacillus cereus* count of *Claris lazera* (n=50).

Bacterial count	Total psychrotrophic	<i>Pseudomonas</i>	<i>Bacillus cereus</i>
Minimum	2.0X10 ³	4.0X10 ²	< 10 ²
Maximum	6.1X10 ⁴	2.0X10 ⁴	3.0X10 ³
Mean	2.5X10 ⁴	6.8X10 ³	8.6X10 ²
Standard error	3.6X10 ³	1.2X10 ³	2.1X10 ²

Table 4: Frequency distribution of the examined *Claris lazera* based on their total psychrotrophic, *Pseudomonas* and *Bacillus cereus* counts (n=50).

Count Range	Total psychrotrophic		Pseudomonas		Bacillus cereus	
	No.	%	No.	%	No.	%
$< 10^2$	25	50	32	64	33	66
$10^2 < 10^3$	0	0	0	0	10	20
$10^3 < 10^4$	7	14	14	28	7	14
$10^4 < 10^5$	18	36	4	8	0	0

$< 10^2$ mean negative result.

DISCUSSION

The psychrotrophic bacteria have been received an increased attention by several investigators during recent years because the modern developments in fish production which resulted in that fish must be held for longer period at low temperature before transportation, processing manufacture or consumption.

Regarding the results recorded in Table (1) it is obvious that the total psychrotrophic count of the examined *Tilapia nilotica* samples had a count ranged from 1.7×10^3 to 3×10^4 with a mean value of $3.2 \times 10^2 \pm 1.3 \times 10^2$, while the total *Pseudomonas* count ranged from 7×10^2 to 2.6×10^4 with a mean value of $4.5 \times 10^3 \pm 1.3 \times 10^3$. Finally, the *Bacillus cereus* count had minimum, maximum and a mean values of 7.8×10^2 , 8×10^3 and $3 \times 10^3 \pm 3.6 \times 10^2$ respectively.

Variable figures were reported by Hassan (1991) who recorded that the log. mean value was 7.8 ± 7.2 . Mousa and Mahmoud (1991) recorded that the mean values of the total psychrotrophic counts of examined *Tilapia nilotica* was $0.56 \times 10^2 \pm 0.028 \times 10^2$ /g. While Mahmoud (1994) recorded that the log. mean values of the total psychrotrophic count in the examined *Tilapia nilotica* was 5.84 ± 0.090 . Higher counts were recorded by Hobbs (1983), Jay (1986) and Hayes (1992). The presence of psychrotrophic organisms in fish may be attributed to contamination from environment around the fish. The processing, equipments, workers, containers, boxes, as well as using polluted water during transportation play an important role for increase counts of psychrotrophic bacteria. Also delaying refrigeration after harvesting and

other handling errors between harvesting and processing lead to decomposition of the fish and allow microbes to grow rapidly (Marriott, 1997).

Pseudomonas species was isolated from fresh water fish by Gram (1993), Mousa and Mahmoud (1997) and Walaa Omer (2004). Lower *Pseudomonas* counts in the examined *Tilapia nilotica* samples was recorded by Lamada-Hanan (1999) with a mean value of $5.4 \times 10^2 \pm 1.1 \times 10^2$. *Pseudomonas* species are widely distributed in nature, polluted water, unsanitized equipments, fishermen hands during harvesting, transportation and storage are the source of fish contamination (Venugopal, 1990).

Bacillus cereus was widely distributed in nature, water, soil, air and dust and can be isolated from a wide variety of foods which it may be normally present. The frequency distribution of the total psychrotrophic count of *Tilapia nilotica* was recorded in table (2) and revealed that the majority of samples lied between 10^4 to 10^5 , while the majority of such examined samples had *Pseudomonas* and *Bacillus cereus* count between 10^3 to 10^4 .

Regarding *Claris lazera* fish samples examined, the result reported in Table (3) revealed that psychrotrophic count ranged from 2×10^3 to 6.1×10^4 with a mean value $2.5 \times 10^4 \pm 3.6 \times 10^3$ /g. While the total *Pseudomonas* count ranged from 4×10^2 to 2×10^4 with a mean value $6.8 \times 10^3 \pm 1.2 \times 10^3$ /g, and finally, the *Bacillus cereus* count ranged from $<10^2$ to 3×10^3 with a mean value of $8.6 \times 10^2 \pm 2.1 \times 10^2$ /g.

Mousa and Mahmoud (1997) found a lower psychrotrophic count in the examined *Claris lazera* fish samples where the mean value was $0.78 \pm 0.022 \times 10^3$.

Table (4) indicated that the frequency distribution of the total psychrotrophic count of the examined *Claris lazera* had a majority count lied between 10^4 to $<10^5$, while *Pseudomonas* and *Bacillus cereus* counts lied between 10^3 to $<10^4$ and 10^2 to $<10^3$ respectively.

Growth of psychrotrophic bacteria in fish has become a significant problem due to the wide spread use of refrigerated storage of fish, and finally we must know that psychrotrophic bacteria when grow in fish can induce different varieties of off-flavour including fruity, putrid, ranced flavour as well as other physical defects.

REFERENCES

- Altayar, M. and Sutherland, A.D. (2006):* Bacillus cereus is a common in the environment but emetic toxin producing isolates are rare. J. Appl. Microbiology, 100(1): 7-14.
- APHA "American Public Health Association" (1984):* Compendium of Methods for Microbiological Examination of Foods. 2nd. Edition. Washington, D.C.
- APHA "American Public Health Association" (1992):* Compendium of Methods for Microbiological Examination of Foods. 3rd Ed. Washington, D.C.
- Berrang, M.E.; Brachett, R.E. and Beuchat, L.R. (1989):* Growth of Listeria monocytogenes on fresh vegetables stored under a controlled atmosphere. J. Food Prot. (52): 702-705.
- Carraseal, A.; Arrienta, G. and Mattar, S. (2002):* Preliminary study of microbiological quality of foods on the Colombian Atlantic Coast. Informe Quincenal-Epidemiologico Nacional, 7(11): 163-169.
- Gram, L. and Melchiorsen, J. (1996):* Interaction between fish spoilage bacteria Pseudomonas spp. and shewanella putrefaciens in fish extracts and on fish tissue. J. Appl. Bacteriol. 80(6): 589.
- Hassan, M.T. (1991):* Microbiological status of frozen dressed fish. M.V.Sc. Thesis Fac. Vet. Med., Moshtohor, Zagazig University-Benha.
- Hayes. P.R. (1992):* Food Microbiology and Hygiene. 2nd ed. El-Sevier Science Publishers (L.Td).
- Hobbs, G. (1983):* Microbial spoilage of fish. In food advances and prospects report. T.A. and Skinner, F.A. (Eds), Society for App. Bact. Symposium series. No. 11, Academic Press London, 217-229.
- ICMSF "International Commission on Microbiological Specification for foods" (1978):* Microorganisms in Foods, Their Significance and Methods of Enumeration. 2nd Ed. Univ. of Toronto Press. Toronto, Buffalo. Canada.
- ISO (1987):* Microbiology-General guidance for enumeration of Bacillus cereus. ISO 7932. Geneva, Switzerland.
- Jay, M.J. (1986):* Modern Food Microbiology. 3rd ed. CBS Publisher and distributors. Indian-Edition.
- Jay, M.J. (2000):* Modern Food Microbiology. Sixth Ed. Gaithersburg. Maryland.

- Lamada-Hana, M.O. (1999):* Chemical and sensory changes associated with microbial flora of Mediterranean boque (Boops boops) stored aerobically at 0, 3, 7 and 10°C. *Appl. Environ. Microbiol.* 65(2): 698.
- Lawsan, T.B. (1970):* Some aspects of fish inspection and public health. *Vet. Reas.*, 87: 528.
- Mahmoud, Y.E. (1994):* Studies on frozen fish ph.D. Thesis. Fac. Vet. Med., Moshtohor. Zagazig University. Benha.
- Marriott, N.G. (1997):* Essential of food sanitation. Consulting Editor, Gill Robertson Chapman and hall 115 5th avenue, New York.
- Mousa, M.M. and Mahmoud, Y.E. (1997):* Hygienic quality of newly caught fresh fish. *Assiut. Vet. Med. J.* 73: 219-233.
- Omer, I.A. Walla (2004):* Antibiotic resistance of *Pseudomonas* species isolated from fish and farm animals. Ph.D. Thesis. Fac. vet. Med. Alexandria.
- Parry, J.M.; Turnbull, P.C.B. and Gibson, J.R. (1983):* A colour Atlas of *Bacillus* species. Wolf Medical Publication Ltd. London.
- Russell, S.M.; Fletcher, D.L. and Cox, N.A. (1995):* Spoilage bacteria of fresh broiler chicken carcasses. *Poultry Sci.*, 74: 2041-2047.
- Venugopal, V. (1990):* Extracellular proteases of contaminant bacteria in fish spoilage. *J. Food Prot.*, 53: 341-350.
- Yumoto, I.; Iwata, H.; Sawabe, T.; Meno, K.; Ichise, N.; Matsuyama, H.; Okuyama, H. and Kawasaki, K. (1999):* Characterization of facultatively Psychrophillic bacterium, *vibrio rumoiensis* sp. Nov., that exhibits high catalase activity. *Appl. Environ. Microbiol.* 65(1): 67-72.