

Dept. of Food Hygiene,  
Faculty of Vet. Med., Assiut University.

## **EFFECT OF NISIN AND CINNAMON ON THE GROWTH AND SURVIVAL OF ENTEROTOXIGENIC METHICILLIN-RESISTANT STAPH. AUREUS IN STERILE MILK AND ICE CREAM**

(With 4 Figures)

By

**S. NASR; NAGAH M. SAAD and RANIA M. EWIDA**

(Received at 19/3/2009)

**تأثير النيسين والقرفة على نمو وبقاء ميكروب المكور العنقودي الذهبى المفرز  
للسموم والمقاوم للميثيسيلين فى اللبن المعقم والآيس كريم**

**سعد الدين نصر ، نجاح محمد سعد ، رانيا محمد عويضة**

تم دراسة تأثير تركيزات مختلفة من النيسين ( 100 ، 200 وحدة/مل) على نمو الميكروب المكور العنقودي الذهبى المفرز للسموم والمقاوم للميثيسيلين المحقون فى عينات اللبن المعقمة معمليا والمضاف إليها ( $10 \times 10^7$  /مل) من الميكروب والذي تم عزله مسبقاً من عينات لبن الخام حيث حفظت العينات عند درجات حرارة الغرفة ( $20 \pm 2^\circ\text{C}$ ) والثلاجة ( $4 \pm 2^\circ\text{C}$ ) وقد تم فحص العينات بعد الحقن مباشرة وكل اثنى عشرة ساعة. وقد أسفرت الدراسة على أن تركيز النيسين ودرجة الحرارة هما العاملان المؤثران على بقاء هذا الميكروب ، ولم يتم عزل الميكروب بعد 72 ساعة فى التركيز ( 200 وحدة/مل) بينما لم يتم عزله بعد 96 ساعة فى التركيز (100 وحدة/مل) من التخزين فى درجة حرارة الغرفة. وعلى الجانب الآخر لم يتم عزل الميكروب بعد 48 و 24 ساعة من التخزين فى درجة حرارة الثلاجة وعند تركيزات (100 ، 200 وحدة/مل) على الترتيب. وقد أشتملت الدراسة أيضا على دراسة تأثير القرفة بنسب 0.3 و 0.6% على الميكروب المكور العنقودي الذهبى المقاوم للميثيسيلين والمفرز للسموم فى عينات الآيس كريم المصنع معمليا والتي تم حفظها تحت درجات تجميد  $-4 \pm 2^\circ\text{C}$  و  $-18 \pm 2^\circ\text{C}$ ، وقد تم فحص العينات مباشرة بعد التصنيع واليوم الأول والثانى والثالث ثم أسبوعياً لمدة 4 أسابيع لعد ميكروب المكور العنقودي الذهبى. ولخصت الدراسة إلى أن درجة حرارة التجميد  $-18 \pm 2^\circ\text{C}$  مع التركيز 0.6% كان لهما دور كبير فى تثبيط نمو الميكروب حيث أنه لم يتم عزله بعد أسبوع واحد من التخزين.

### **SUMMARY**

The effect of nisin on the survival of enterotoxingenic methicillin-resistant *Staph. aureus* (MRSA) was evaluated using different concentrations of nisin (0.00, 100 and 200 IU/ml) in a laboratory

prepared sterile milk and inoculated with the previously isolated and identified MRSA to yield a concentration of  $1 \times 10^7$  cfu/ml. The inoculated sterile milks were kept at room temperature ( $20 \pm 2^\circ\text{C}$ ) and refrigerator temperature ( $4 \pm 2^\circ\text{C}$ ). MRSA counts were determined using Oxacillin Resistance Screen Agar Base (ORSAB) supplement with two antibiotics-oxacillin at (2 mg/L) and polymyxin B (50.000 IU/L). ORSAB and pH value were determined every twelve hours. MRSA strains couldn't be detected after 96 and 72 h in the samples of sterile milk containing nisin in concentrations of 100 and 200 IU/ml, while, in the control sample the MRSA survived till the end of the 96 h of storage at room temperature. Moreover, MRSA failed to be detected after 48 and 24 h in the samples of sterile milk containing nisin in concentrations of 100 and 200 IU/ml and stored at refrigerator temperature, respectively. Ice cream was prepared at the laboratory to study the effect of ground cinnamon in concentrations of 0.3 and 0.6% on the growth and survival of enterotoxigenic MRSA a concentration of  $2 \times 10^7$  cfu/ml at freezing ( $-4 \pm 2^\circ\text{C}$ ) and deep freezing ( $-18 \pm 2^\circ\text{C}$ ) temperatures. The obtained results showed that the advantage of using 0.6% is better than using of 0.3% of ground cinnamon.

**Key words:** Enterotoxigenic MRSA, milk, nisin, ice cream, cinnamon.

## INTRODUCTION

MRSA infections are of the global health issues due to the severity of the illnesses that may occur. The roles and the sources of food contamination are still unclear since only few reports on the presence and possible origin of MRSA in foods are available. In his analysis of specimens from food-producing animals, including milk, Lee (2003) found 15 strains harbouring the *mecA* gene and most of the MRSA isolates were from milk. He concluded that contaminated foods of animal origin may represent a source of MRSA infection for humans. Kitai *et al.* (2005) suggested that food handlers were the source of contamination. This finding emphasized the role of humans as an important reservoir of MRSA. In the two foodborne acquired MRSA outbreaks, food handlers carrying the epidemic strains in their nostrils were identified to be the source of contamination of the incriminated food (Kluytmans *et al.*, 1995 and Jones *et al.*, 2002).

Food additive is a substance or mixture of substances, other than the basic food stuff, which are present in food as a result of any aspect of production, processing, storage, or packaging. Consequently,

biopreservatives and natural preservatives are needed which possess antimicrobial activity and cause no problems to the handlers and consumers. In this respect, nisin and cinnamon were tested for their inhibitory activity towards the growth of some microorganisms and used as food additives.

Nisin is a natural antimicrobial peptide of 34 amino acids produced by *Lactococcus lactis subsp. lactis* (Carr *et al.*, 2002). The peptide is suggested to be effective against a wide range of Gram positive bacteria including *Staph. aureus* (Narasimhan *et al.*, 1988 and Choi *et al.*, 2000). In the U.S., nisin was confirmed to be "generally recognized as safe" in 1988 (F.D.A., 1988). Because of its high antibacterial activity and nontoxicity for humans, nisin has already been employed as food preservative for a long time and is licensed by 48 countries around the world (Deegan *et al.*, 2006).

Cinnamon has been used since ancient times both as a culinary spice and for medicinal and other purposes. The medical properties of cinnamon were utilized by ancient Egyptians. Its mild anti-inflammatory, anti-spasmodic, and anti-clotting properties are believed to be due to its content of cinnamaldehyde. Cinnamon extracts have also inhibited the growth of cultured tumor cells (Craig, 2008). Moreover, cinnamon can be useful in the treatment of type 2 diabetes; as well as lowering triglyceride levels and serum cholesterol (Khan *et al.*, 2003). Also, it is useful as a food preservative to inhibit the growth of common foodborne bacteria as *Staph. aureus* and MRSA (Chang *et al.*, 2001; Yuste and Fung, 2003; Prabuseenivasan *et al.*, 2006 and Ağaoğlu *et al.*, 2007).

This work was undertaken to throw light on the effect of adding different concentrations of nisin and cinnamon on viability of enterotoxigenic MRSA in sterile milk and ice cream.

## **MATERIALS and METHODS**

### **I- Effect of different concentrations of nisin on the growth and survival of enterotoxigenic MRSA in laboratory prepared sterile milk:**

#### **- The organism:**

Enterotoxigenic MRSA strain used was previously isolated and well identified from the examined milk samples.

**- Experimental procedure:**

Raw milk was laboratory sterilized by autoclaving. The sterile milk inoculated with MRSA in concentration  $1 \times 10^7$  cfu/ml and nisin was added in concentration of 0.00 (nisin-free), 100 and 200 IU/ml. Each sample was divided into two portions, the first portion was kept at room temperature ( $20 \pm 2^\circ\text{C}$ ) and the other was stored at refrigerator temperature ( $4 \pm 2^\circ\text{C}$ ). The effect of nisin on the growth and survival of MRSA was determined by using ORSAB after 3 h and then every 12 h and compared with nisin-free samples.

**- Measurement of pH value:**

The pH value of each sample was determined according to standard methods of A.P.H.A. (1992) with a pH meter (Orion Research model 3200 A/digital analyzer) previously standardized with buffer solutions of pH 3.0 and pH 7.0 buffer solutions.

**II- Effect of different concentrations of ground cinnamon on the growth and survival of enterotoxigenic MRSA in laboratory made ice cream:**

**- Organism:**

Enterotoxigenic MRSA strain used was previously isolated and well identified from the examined ice cream samples.

**- Experimental procedure:**

Ice cream samples were prepared in the laboratory according to the manufacture (Egyptian Dairy & Food Company) and then the mixture inoculated with MRSA culture to contain  $2 \times 10^7$  cfu/ml. After that, the ground cinnamon was added during manufacture of these samples in concentrations of 0.00 (cinnamon-free), 0.3 and 0.6%. Each sample was divided into two portions, the first portion was kept at freezing temperature ( $-4 \pm 2^\circ\text{C}$ ) and the other was stored at deep freezing temperature ( $-18 \pm 2^\circ\text{C}$ ).

Samples from the two portions were taken before and after addition of ground cinnamon to determine the initial count of the organism by using ORSAB, the count after hardening of ice cream samples, after the first, second and third day. Then, the samples were tested every week up to 4 weeks of storage for MRSA count.

## **RESULTS**

The obtained results were recorded in Figures 1-4.

**Fig. 1:** Effect of different concentration of added nisin on the growth and survival of MRSA in sterile milk stored at room temperature ( $20\pm 2^{\circ}\text{C}$ ).

**Fig. 2:** Effect of different concentration of added nisin on the growth and survival of MRSA in sterile milk stored at refrigerator temperature ( $4\pm 2^{\circ}\text{C}$ ).

**Fig. 3:** Effect of different concentration of added ground cinnamon on the growth and survival of MRSA in laboratory made ice cream stored at freezing temperature ( $-4\pm 2^{\circ}\text{C}$ ).

**Fig. 4:** Effect of different concentration of added ground cinnamon on the growth and survival of MRSA in laboratory made ice cream stored at deep freezing temperature ( $-18\pm 2^{\circ}\text{C}$ ).

## DISCUSSION

### **I- Effect of different concentrations of nisin on the growth and survival of enterotoxigenic MRSA in laboratory prepared sterile milk:**

The data presented in Fig. 1 showed that the count of MRSA in sterilized milk stored at room temperature containing nisin 100 IU/ml was increased to  $4.0 \times 10^8$  cfu/ml at the 3<sup>rd</sup> h of the storage, then it reached  $3.0 \times 10^7$  at the 12<sup>th</sup> h of the storage. Then at the end of first 24 h of the storage the count gradually decreased and reached  $8.5 \times 10^5$  cfu/ml. The count of MRSA decreased until the pathogen failed to be detected on the plate and could not be isolated at the end of 96 h. While, sterile milk containing nisin 200IU/ml, the count of MRSA increased  $1.4 \times 10^8$  cfu/ml after 3 h of storage. The count gradually decreased until the pathogen failed to be detected on the plate at the end of 72 h of the storage and failed to be isolated from the milk. On the other side, in control sample, MRSA increased from the initial count of  $1 \times 10^7$  to  $4.5 \times 10^9$  cfu/ml in 96 h. Gradual decrease of pH value of the sterile milk occurred from 6.40 to 5.62 in 0 time to the end of 96 h, respectively.

Result in Fig. 2 depicted the effect of nisin on the survival of MRSA in sterile milk stored at refrigerator temperature ( $4 \pm 2^\circ\text{C}$ ). The initial count of MRSA was  $1.0 \times 10^7$  cfu/ml decreased to  $2.0 \times 10^6$  and  $3.0 \times 10^5$  cfu/ml after 3<sup>rd</sup> h after storage 100 and 200 IU/ml nisin containing samples, respectively. The organism failed to be detected on the plate and could not be isolated from the samples containing 100 and 200 IU/ml nisin after 48 and 24 h storage at ( $4 \pm 2^\circ\text{C}$ ), respectively. While, the pathogen survived in the sample free from nisin up to 48 h of the storage at refrigerator temperature. Slight decrease of pH of the sterile milk from 6.40 to 6.06 at the end of 48 h of the storage.

From the aforementioned results, it was evident that the increased concentration of nisin and the storage of the sterile milk at refrigerator temperature had an inhibitory effect on the survival of *Staph. aureus*. These results agree with the theory which pointed out that *Staph. aureus* is a robust bacterium and can survive for long period at low temperatures below those which permit growth. Yet, refrigeration at  $<4^\circ\text{C}$  may be considered the only viable method for control of growth and toxins production (I.C.M.S.F., 1996 and Ryser, 2001).

Also, the obtained results agree with those recorded by Rilla *et al.* (2004), who found the inhibitory effect of *Lactococcus lactis subsp. lactis* IPLA 729 on MRSA inoculated into milk in different inoculums ( $1.8 \times 10^4$

and  $7.2 \times 10^6$  cfu/ml). While, MRSA when growing alone, it reached levels of  $6.4 \times 10^8$  and  $1.9 \times 10^9$  cfu/ml, respectively. In both cases, a slight growth of MRSA occurred within the first 3 h of incubation. However, complete disappearance of the pathogen occurred at the end of the incubation period.

Many investigators recorded the inhibitory effect of nisin on *Staph. aureus* as Aman and Ahmed (1997), they added nisin in concentration of 50, 100, 150, and 200 IU/ml of nisin in cheese milk had a highly bactericidal effect against *Staph. aureus*. Also, Narasimhan *et al.* (1988) and Choi *et al.* (2000) reported the inhibitory effect of nisin on *Staph. aureus*.

Numerous studies have been made on the mode of action of nisin on susceptible vegetative cells. The target of the inhibition effect of nisin is the cytoplasmic membrane where it depolarizes energized bacterial membranes (reduces transmembrane potential) and form voltage-dependent multistate pores. The result of a pore formation is the loss of accumulated amino acids and the inhibition of amino acid transport (Shehata, 1981). Delves (1990) indicated that nisin causes disruption, either resulting in leakage of essential cellular material such as adenosine triphosphate from the cell or in more severe cases lysis of the cell.

## **II- Effect of different concentrations of ground cinnamon on the growth and survival of enterotoxigenic MRSA in laboratory made ice cream:**

In ice cream samples stored at freezing temperature ( $-4 \pm 2^\circ\text{C}$ ) and containing 0.3% ground cinnamon (Fig. 3), the counts of MRSA diminished gradually from  $1.1 \times 10^4$  to  $2.8 \times 10^2$  cfu/ml by the end of 3<sup>rd</sup> week, and failed to recover from the samples by the end of the 4<sup>th</sup> week. The ice cream samples containing 0.6% ground cinnamon, the counts of MRSA decreased gradually from  $8.0 \times 10^3$  to  $3.0 \times 10^2$  cfu/ml by the end of the 2<sup>nd</sup> week, and failed to be detected in the samples by the end of 3<sup>rd</sup> week. Concerning cinnamon free ice cream, the number of MRSA decreased gradually during the storage period to reach its minimum level  $6.0 \times 10^4$  cfu/ml by the end of the 4<sup>th</sup> week.

As presented in Fig. 4, the numbers of MRSA ( $4.7 \times 10^3$  and  $2.1 \times 10^3$  cfu/ml) reduced in ice cream samples with 0.3 and 0.6% added ground cinnamon during the weeks of storage at the deep freezing temperature ( $-18 \pm 2^\circ\text{C}$ ). No viable MRSA could be detected in the samples by the end of 3<sup>rd</sup> and 1<sup>st</sup> week, respectively.



These findings were in agreement with the theory that cinnamon had antibacterial activities against many of bacteria including methicillin resistant *Staph. aureus* (Chang *et al.*, 2001). Also, many other investigators reported the inhibitory effect of cinnamon against *Staph. aureus* as Yuste and Fung (2003), Prabuseenivasan *et al.* (2006) and Ağaoğlu *et al.* (2007).

## REFERENCES

- A.P.H.A. (1992): Standard Methods for the Examination of Dairy Products. 16<sup>th</sup> Ed., American Public Health Association. New York.
- Ağaoğlu, S.; Dostbil, N. and Alemdar, S. (2007): Antimicrobial activity of some spices used in the meat industry. Bull. Vet. Inst. Pulawy, 51: 53-57.
- Aman, I.M. and Ahmed, H.F. (1997): Incidence and survival of some foodborne pathogens in milk and cheese. J. Egypt. Vet. Med. Assoc., 57(1): 151-163.
- Carr, F.J.; Chill, D. and Maida, N. (2002): The lactic acid bacteria: a literature survey. Crit. Rev. Microbiol., 28: 281-370.
- Chang, S.; Chen, P. and Chang, S. (2001): Antibacterial activity of leaf essential oils and their constituents from *Cinnamomum osmophoeum*. J. Ethnopharmacology, 77: 123-127.
- Choi, H.J.; Cheigh, C.I.; Kim, S.B. and Pyun, Y.R. (2000): Production of a nisin like bacteriocin by *Lactococcus lactis subsp. lactis* A164 isolated from kimchi. J. Appl. Microbiol., 88(4): 563-571.
- Craig, W. (2008): Cinnamon. [http://www.vegetarian-nutrition. Info/ herbs/ cinnamon. php](http://www.vegetarian-nutrition.Info/herbs/cinnamon.php).
- Deegan, L.H.; Cotter, P.D.; Hill, C. and Ross, P. (2006): Bacteriocins: biological tools for bio-preservation and shelf-life extension. Int. Dairy J., 16: 1058-1071.
- Delves, B.J. (1990): Nisin and it's uses as a food preservative. Food Technol., 11: 100-113.
- F.D.A. (Food and Drug Adminstration) (1988): Nisin preparation: affirmation of GRAS status as direct human food ingredient. Fed. Regist., 53: 11247-11251.

- I.C.M.S.F. (*International Commission on Microbiological Specification*) (1996): International Committee on Microbiological Specification of Foods. Microorganisms in Foods, Microbiological Specifications of Food Pathogens. 1<sup>st</sup> Ed., Chapman and Hall, 2-6 Boundary Row, London SE1 8 HN, UK.
- Jones, T.F.; Kellum, M.E.; Porter, Susan; Bell, M. and Schaffner, W. (2002): An outbreak of community-acquired foodborne illness caused by Methicillin resistant *Staph. aureus*. *Emerging Infectious Dis.*, 8(1): 8-11.
- Khan, A.; Safdar, M.; Ali Khan, M.M.; Khattak, K.N. and Anderson, R.A. (2003): Cinnamon improves glucose and lipids of people with type 2 diabetes. *Diabetes Care*, 26(12): 3215-3218.
- Kitai, S.; Shimizu, A.; Kawano, J.; Sato, E.; Nakano, C.; Uji, T. and Kitagawa, H. (2005): Characterization of methicillin resistant *Staph. aureus* isolated from retail raw chicken meat in Japan. *J. Vet. Med. Sci.*, 67: 107-110.
- Kluytmans, J.; Leeuwen, W.V.; Goessens, W.; Hollis, R.; Hawn Messer, S.; Herwaldt, L.; Bruining, H.; Heck, M.; Rost, J.; Leeuwen, N.V.; Belkum, A.V. and Verbrugh, H. (1995): Food-initiated outbreak of methicillin resistant *Staph. aureus* analyzed by pheno- and genotyping. *J. Clin. Microbiol.*, 33(5): 1121-1128.
- Lee, J. (2003): Methicillin (Oxacillin) resistant *Staph. aureus* strains isolated from major food animals and their potential transmission to humans. *Appl. Environ. Microbiol.*, 69(11): 6489-6494.
- Narasimhan, R.; Padmanaban, V.D. and Ulaganathan, V. (1988): Effect of sodium propionate on the growth profile of microorganisms (antimicrobial activity of sodium propionate). *Cheiron.*, 17(4): 144-150.
- Prabuseenivasan, S.; Jayakumar, M. and Ignacimuthu, S. (2006): In vitro antibacterial activity of some plant essential oils. *BMC Complement Altern. Med.*, 6: 39.
- Rilla, Natalia; Martinez, B. and Rodriguez, Ana (2004): Inhibition of a methicillin resistant *Staph. aureus* strain in Afueg'l Pitu cheese by the nisin Z-producing strain *Lactococcus lactis subsp. lactis* IPLA 729. *J. Food Prot.*, 67(5): 928-933.
- Ryser, E.T. (2001): Public health concerns. In: *Applied Dairy Microbiology*. Marth, E.H. and Steele, J.L. (eds.). Marcel Dekker, Inc., New York, pp.: 478-486.

- Shehata, T.E. (1981): Effect of selected food additives on the growth of psychrophilic strains of *Bacillus*. II Effect of nisin, sodium benzoate and potassium sorbate on growth inhibition of *B. cereus* in nutrient broth and in skim milk. Research-Bulletin, 1554: 12.
- Yuste, J. and Fung, D.Y.C. (2003): Evaluation of *Salmonella typhimurium*, *Yersinia enterocolitica* and *Staph. aureus* count in apple juice with cinnamon, by conventional media and thin agar layer method. Food Microbiol., 20: 365-370.




Figure 1. Effect of different concentration of added nisin on the growth and survival of MRSA in sterile milk stored at room temperature ( $20\pm 2^{\circ}\text{C}$ ).




Figure 2. Effect of different concentration of added nisin on the

growth and survival of MRSA in sterile milk stored at refrigerator temperature ( $4\pm 2^{\circ}\text{C}$ ).




Figure 3. Effect of different concentration of added ground cinnamon on the growth and survival of MRSA in laboratory made ice cream stored at freezing temperature ( $-4\pm 2^{\circ}\text{C}$ ).




Figure 4. Effect of different concentration of added ground cinnamon on the growth and survival of MRSA in laboratory made ice cream stored at deep freezing temperature ( $-18\pm 2^{\circ}\text{C}$ ).