Incidence of campylobacter species in milk and some milk products

By

Nagah Saad; Ahmed, A-H, Ahmed; Amal Ali Abdel – Haleem *Thanaa Nassife* *Animal Health research Institute, Assiut.

<u>Summary</u>

Three hundred and fifty random samples of raw milk (150) and some milk products including kareish and Damietta cheese, ice-cream and cooking butter (50 samples each) were collected from Assiut city Markets, dairy shops, and dairy farms. The samples were examined for isolation and identification of Campylobacter spp.

The obtained results revealed that 10 (6.7%), 7 (14%), 5 (10%) of the examined raw milk, kareish cheese and ice cream samples were contaminated by Campylobacter spp. using Brucella agar medium. However, the incidence of Campylobacter spp. Using Campylobacter agar was 9 (6%) in raw milk samples, 3 (6%) in kareish cheese, 6 (6%) in ice cream, 1 (2%) in Damietta cheese and 3 (6%) in cooking butter. The isolated Campylobacter spp. could be identified as Campylobacter jejuni, C coli, C laridis, C. fetus, C. hyointestinalis and C. fecalis.

Plasmid profile and antibiogram of the isolated Campylobacter jejuni recovered from the examined raw milk and dairy products revealed that 5 out of 10 isolates (50%) of C. jejuni carry (1-2) plasmids of high molecular weight with resistance to Cephalothin, Oxytetracycline, Flemkuin and Kanamycin and sensitivity to Norfloxacin, Enrofloaxacin, Gentamycin and Nalidixic acid. The bublic heath significance and suggestive measures to improve the keeping quality as well as sanitary conditions of milk and milk products were given.

Introduction

In the last few years Campylobacters have emerged as the most frequent cause of acute bacterial gastroenteritis in man (Casini et al., 1997) as the number of human cases of Campylobacteriosis has increased dramatically in recent years in many countries (Nielsen et al., 2000).

The term Campylobacter is a Greek name composed of campy, means curved and bacter, means rod i.e curved rods which describes the appearance of the organisms (Nachamkin et al., 1992).

The genus Campylobacter comprises many species (14 species, subspecies, biovars, with 17 official names) (AbdeI-Samei, 2000), which are newly established as bacterial agents of clinical importance in humans. The most frequently identified human pathogenic species of Campylobacter are C. jejuni and C. coli which are closely related and their infections appear to share many clinical and epidemiologic characteristic in addition to C. laridis. They account for more than 99% of the human isolates of infection (C. jejuni 90%). Recent work suggests that C. upsaliensis is also enteropathogenic, and occasionally other species such as C.

Key words: Campylobacter spp, milk, milk products, plasmid, antibiotic sensitivity.

hyointestinalis was isolated from patients with diarrhea. (Fennell et al., 1984; Edmonds et al., 1987 and Skirrow, 1990).

Campylobacter may found as a normal intestinal flora of both wild and domesticated animals specially those used for food production (Penner, 1988). However, the main source of Campylobacter infection is probably raw milk and milk products which are the most commonly implicated vehicles in food - borne outbreaks of Campylobacter enteritis. (Richter et al., 1992 and Bean et al., 1996). Contamination of milk can occur by direct excretion from an asymptomatic cow with mastitis (Hutchinson et al., 1985) or through fecal contamination during milking from cattle infected or colonized with the organism (Waterman et al., 1984 and Humphrey and Beckett 1987). Also post pasteurization contamination of milk and dairy products have been found responsible for Campylobacter outbreaks (White, 1986).

The mechanism of Campylobacter pathogenesis has not been elucidated fully, as several virulence related factors such as adherence, invasiveness and production of heat labile cholera-like enterotoxin or cytotoxin have been associated with the organism (Walker et al., 1986 and Kalman et al, 2000). C. jejuni has become the most commonly reported cause of food-bome enteritis in people world wide (Skirrow, 1990 and Butzler and Oosterom, 1991). It can produce different types of toxins as enterotoxin, cytolethal distending toxin and cytolethal rounding toxin.

The infective dose of C. jejuni ranges from 500 to 10.000 cells, (Doyle, 1991; Reed, 1994 and Phillips, 1995). Campylobacter enteritis has been associated with some complications such as arthritis, recurrent colitis, Heamolytic Uremic Syndrome (HUS), Reiter syndrome, a reactive arthropathy which develops in about 1% of patients 1-2 weeks after onset of illness, Miller-fisher syndrome, Chinese paralytic syndrome, Guillian-Barre syndrome (GBS) particularly in immunocompromized individuals, which is, a disorder resulting in acute neuromuscular paralysis, as a serious sequelae of Campylobacter infection, up to 40% of patients with GBS have evidence of recent Campylobacter infection (Smith, 1995 and Allos, 1997).

Although Campylobacter doesn't commonly cause death, it has been estimated that approximately 200 persons with Campylobacter infections may die each year in the United States (Skirrow 1990 and Tauxe 1992). The vast majority of outbreaks of Campylobacteriosis have been associated with consumption of unpasteurized or inadequately pasteurized cow's milk in New Zealand, Scotland, Switzerland and England (Hutchinson et al., 1985; Hudson et al., 1990) as well as dairy products (Barrett, 1986). In Egypt, acute diarrhea of presumed infectious origin is responsible for more than 50% of deaths for those under two years of age (Ewyda, 1990).

Therefore, Campylobacter spp. continue to be highly important human pathogens and there is an increase interest of Campylobacters particularly C. jejuni as a health risk affecting both human and animal and because of the involvement of milk and milk products in human Campylobacter enteritis, the present work was planned to study the following items:

- Occurrence of Campylobacter spp. in milk and some milk products.
- Identification of the isolated Campylobacter spp.
- Plasmid profile and antibiotic sensitivity of the isolated strains of C. jejuni.

MA TERIAL AND METHODS

1. Isolation of Campylobacter spp. from milk and some milk products:

1. Collection of samples:

A total of 350 random samples of raw milk (150), (Damietta and kareish cheese), ice-cream and cooking butter (50 samples each) were purchased from different localities in Assiut city. The samples were collected in clean, dry and sterile containers while, ice cream-samples were taken in an ice-box. Collected samples were transferred to the laboratory as soon as possible to be examined. Each milk sample was mixed by inversion several times and tested for heat treatment using Storch's test (Lampert, 1975).

<u>2. Preparation of samples:</u> The samples were prepared according to the technique recommended by (A.P.H.A., 1992).

3. Experimental procedures: The technique adopted by Boer et al. (1984) was used.

3.1.1- Enrichment procedure:

1 ml of each prepared sample was inoculated into Campylobacter enrichment broth and brucella broth. Each broth containing 5% lysed horse blood, Skirrow Campylobacter selective supplement and Skirrow Campylobacter growth supplement. The inoculated tubes were incubated at 42°C for 24 h in an atmosphere of 5% oxygen, 10% carbon dioxide and 85% nitrogen using an anaerobic jar and Campylobacter gas generating kits (Oxoid, 1990).

3.1. 2- Selective plating:

Incubated broth cultures were then streaked onto plates of both Brucella and Campylobacter blood agar base supplemented with Skirrow Campylobacter selective supplement, 5% lysed horse blood, and Skirrow Campylobacter growth supplement. Streaked plates were incubated at 42°C for 48 h under appropriate microaerophilic conditions in anaerobic jar with activated gas generating kit by using the Gas pack system BBL.

All pure cultured colonies were subjected to identification scheme according to Skirrow (1990), Rosef and Yundestad (1982) and Smith et al. (1997).

II. Plasmid DNA Extraction:

It has been carried out in the Molecular Biology and Genetic Engineering Research Center in Assiut University. Detection of plasmid DNA was done by Agarose Gel Electrophoresis; (Kaufman et al., 1995).

III. Antimicrobial sensitivity testing of Campylobacter jejuni: Using disc diffusion method (Baron et al, 1994)

Results

Table 1	<i>l</i> .	Incidence	of	<i>campylobacter</i>	spp.	in	the	examined	samples	of	milk	and	<u>milk</u>
product	ts.												

		Positive samples								
Examined samples	No. of samples	Brucella a	gar	Campylobacter agar						
		No.	%	No.	%					
Milk	150	10	6.7	9	6					
Damietta cheese	50	-	-	1	2					
Kareish cheese	50	7	14	3	6					
Ice – cream	50	5	10	3	6					
Butter	50	-	-	3	6					
Total	350	22	6.3	19	5.4 %					

Table 2. Incidence of C. jejuni in the examined samples of milk and milk products.

		Positive samples							
Examined samples	No. of samples	Brucel	la agar	Campylobacter agar					
		No.	%	No.	%				
Milk	150	2	1.3	5	3.3				
Damietta cheese	50	-	-	1	2				
Kareish cheese	50	1	2	1	2				
Ice – cream	50	1	2	3	6				
Butter	50	-	-	-	-				
Total	350	4	1.1 %	10	2.9 %				

	No. of identified													
Examined samples	No. of	Total isolates	c. je	juni	С.	coli	C. f	etus	C. laridis C. hyointestinali		yointestinalis	C. fecalis		
	sampies	isouies	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Milk	150	10	2	1.3	2	1.3	2	1.3	3	2	1	0.07	-	-
Damietta cheese	50	-	-	-	-	-	-	-	-	-	-	-	-	-
Kareish cheese	50	7	1	2	3	6	-	-	-	-	1	2	2	4
Ice – cream	50	5	1	2	3	6	-	-	-	-	1	2	-	-
Butter	50	-	-	-	-	-	-	-	-	-	-	-	-	-

Table 3. Incidence of campylobacter spp. in the examined milk and milk products samples using Brucella agar medium.

Table 4. Incidence of campylobacter spp. in the examined milk and milk products samples using Campylobacter agar medium.

								No. of id	entified i	solates				
Examined samples	No. of	Total isolates	c. jejuni		C. coli		C. fetus		C. laridis		C. hyointestinalis		C. fecalis	
	sampies	isolules	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Milk	150	9	5	3.3	4	2.7	-	-	-	-	-	-	-	-
Damietta cheese	50	1	1	2	-	-	-	-	-	-	-	-	-	-
Kareish cheese	50	3	1	2	2	4	-	-	-	-	-	-	-	-
Ice – cream	50	3	3	6	-	-	-	-	-	-	-	-	-	-
Butter	50	3	-	-	2	4	-	-	1	2	-	-	-	-

		Plasmid b	earing isolates	Non plasmid bearing isolates			
Sources of isolates	No. of isolates	No. / 10	%	No. / 10	%		
Milk	5	2	20	3	30		
Damietta cheese	1	1	10	-	-		
Kareish cheese	1	1	10	-	-		
Ice – cream	3	1	10	2	20		
Total	10	5	50%	5	50%		

Table 5. Correlation between C. jejuni isolates and plasmids

Table 6. Antibiotic sensitivity of Compylobacter jejuni isolated from the examined raw milk and milk products samples

Types of antibiotics	Degree of sensitivity of campylobacter jejuni isolates	% of sensitivity
Amoxicillin 30 mg	Moderately sensitive (++)	80%
Cephalothin 30 mg	Resistant	0
Enrofloxacin 5 mg	Moderately sensitive (++)	100%
Flemkuin 30 mg	Weakly sensitive (+)	40%
Gentamycin 10 mg	Moderately sensitive (++)	100%
Kanamycin 30 mg	Weakly sensitive (+)	80%
Naldixic acid 30 mg	Sensitive (+++)	100%
Norfloxacin 10 mg	Sensitive (+++)	100%
Oxytetracycline 30 mg	Weakly sensitive (+)	60%



Fig 1: A garose gel (0.7%) stained with ethidium bromide 0.05% showing:

M : Hundred base pair ladder marker (Sigma)

Lanes 2.7 and 8 : Two copy plasmids of high molecular weight.

Lanes 1 and 6 : single copy plasmid of high molecular weight.

Lanes 3, 4, 5, 9 and 10 showing negative plasmid bearing

Discussion

<u>1- Incidence of Campylobacter spp. in milk and some milk products:</u>

1) Raw milk:

Results recorded in Table 1 showed that Campylobacter spp. were isolated from 10 (6.7%) of 150 examined raw milk samples on Brucella agar and from 9 samples (6%) on Campylobacter blood agar.

These results are nearly in agreement with those obtained by Wegmuiler et al. (1993) and Abdel-Hady (1996). Lower percentages were stated by Inokova and Ivanova (1996); Hudson et al. (1999) and DuzGun et al. (2000). Whereas, higher percentages were recorded by El-Nokrashy et al. (1997); El-Prince et al. (1998) and Roshdy (2000). However, several investigators failed to detect Campylobacter spp. in the examined milk samples such as Mouffok and Lebres (1992) and Federighi et al. (1999).

The variation in these results may be attributed to different factors, including: level of contamination and methods of isolation, variety of enrichment broth systems, high sensitivity of the organism to normal atmospheric concentration of oxygen and to adverse conditions resulting from acid development in raw milk that represent stress factor on the organism resulting in failure of cultural trials even from contaminated samples. (Ray and Johnson, 1984).

According to the results presented in Tables 2 & 3 it is evident that different Campylobacter organisms could be isolated in variant percentages from the examined raw milk samples, these organisms 10 isolates recovered on Brucella agar were identified as C. jejuni (2 isolates), C coli (2 isolates), C. laridis (3

isolates), C. fetus (2 isolates) and C. hyointestinalis (1 isolate). While, on Campylobacter agar, 9 isolates were identified as C. jejuni (5 isolates) and C. coli (4 isolates).

The presence of Campylobacter spp. in the examined raw milk samples may be attributed to poor hygienic conditions under which raw milk is produced and handled, contamination of milk during or after milking is probably of fecal origin, however improper washing and treatment of the udder with suitable disinfectant or contact of the milking pails with the floor may result in a high level of contamination. Furthermore, naturally occurring Campylobacter mastitis and contaminated water supply may act as a source of milk contamination (Mentzing, 1981).

2) Damietta cheese:

As recorded in Tables 1 & 3 it is apparent that out of 50 examined using Campylobacter agar samples of Damietta cheese, one sample was positive for Campylobacter spp. represented by C. jejuni (2%). Using Campylobacter agar This result was in agreement with that (2%) obtained by EI-Nokrashy et al. (1998). Several investigators failed to isolate Campylobacter spp. from the examined cheese samples (Ehlers et al., 1982, Bachmann, 1994 and Federighi et al., 1999).

Lower incidence or absence of the organism in a such product may be due to its fragile nature and sensitivity to adverse conditions of acid development as Campylobacters are inactivated at pH 4.5 (Doyle and Roman, 1981). Water activity and presence of salt (5-15%) which represent stress factors on the organism oftenly result in failure to culture it even from contaminated products or to be recorded as low incidence (Ray and Johnson, 1984). Moreover, Frazier and Westhoff (1988) reported that salt causes high osmotic pressure and hence plasmolysis of cells, dehydrates cheese by drawing out and tying up moisture as it dehydrates microbial cell and it ionizes to yield the chlorine ion, which is harmful to the organism and it interferes with the action of proteolytic enzymes.

3) Kareish cheese:

The results in Table 1 revealed that Campylobacter spp. Were isolated from 7 (14%) of 50 examined kareish cheese samples. According to the results represented in Table 3, it is evident that different Campylobacter organisms could be isolated in variant percentages from the examined kareish cheese samples. These organisms (7 isolates) were identified as C.jejuni (1 isolate) (2%), C. coli (3 isolates) (6%), C. fecalis (2 isolates) (4%) and C. hyointestinalis (1 isolate) (2%) on Brucella agar, while, on Campylobacter blood agar, Campylobacter spp. was isolated from 3 (6%) of the examined kareish cheese samples, these organisms were identified as 1 (2%) C. jejuni and 2 (4%) C. coli as recorded in Table 4. Campylobacter spp. failed to recover from Kareish cheese samples examined by AbdeI-Hady (1993) and Federighi et al. (1999).

As Campylobacter can remain viable in fresh cheese for only a short period of time (Butzler and Oosterom, 1991). The fact that these cheeses may be consumed immediately after production and may pose a public health risk. The relatively high results of this study could be attributed to the neglected sanitary control adopted during manufacturing, handling and distribution of kareish cheese.

4) Ice-cream:

The data presented in Tables 1 & 3 showed that out of 50 examined samples of ice-cream, 5 samples (10%) contained Campylobacter spp, on Brucella agar, these spp. could be identified as 1 (2%) C. jejuni, 3 (6%) C. coli and 1 (2%) C. hyointestinalis. Whereas, on Campylobacter agar Campylobacter spp. was isolated from 3 (6%) of the examined samples represented by C. jejuni. (Table 4) Several investigators failed to isolate Campylobacter spp. from ice cream (Ehlers et al., 1982; Ray and Johnson, (1984) and AbdeI-Hady, 1993). The lower incidence of the organism in such products could be attributed to its sensitivity to

conditions of freezing which stressed Campylobacter organisms and resulted in failure to recover the organism from contaminated frozen food.

5) Butter:

Regarding the results in Tables 1 & 3 out of 50 examined samples of cooking butter 3 (6%) were contaminated with Campylobacter spp. These spp. were identified as 2 (4%) C. coli and 1 (2%) C. laridis. The presence of Campylobacter spp. using Campylobcter in examined cooking butter samples may be due to poor hygienic conditions under which butter is produced and handled. Such contamination is probably of fecal origin.

IV- Plasmid profile and antibiotic susceptibility of campylobacter jejuni:

Bacterial plasmids are extrachromosomal DNA known to be code for toxin production, adhesiveness, antibiotic resistance and serum resistance (Baroun and Ou, 1991 and Lax et al., 1995).

Plasmid analysis was performed in the present study on C jejuni isolated from raw milk and milk products (Damietta cheese, kareish cheese and ice-cream) as well as their sensitivity to some selected antibiotics. The plasmid pattern in Table 5 of the examined strains belonging to C. jejuni showed that 5 (50%) out of the 10 strains bear plasmids of high molecular weight (over 2.6 Kpb) 3 of the 5 isolate carry 2 copies of plasmid as showed in Fig.1, higher percentages were detected by Ansary and Veloo (1991) (62.2%) and Jay (1996) (64.7%). Lower percentages were detected by Lekowska Kochanial et al. (1996) (36%), however, Kalman et al. (2000) failed to detect plasmid DNA of C. jejuni isolate .

The relation between possession of plasmid DNA and the tested isolates and the antimicrobial resistance pattern showed that all the five strains that have plasmid showed a resistance against Cephalothin. Out of 5 isolates, 2 strains showed resistance against Flemukin and 3 out of 5 isolates showed resistance to Oxytetracycline. Table 6 summarized the antibiotic sensitivity of isolated Campylobacter jejuni. All the tested isolates were 100% sensitive to Gentamycin 10 mg, Nalidixic acid 30mg, Norfloxacin 10 mg, Enrofloxacin 5 mg. None of the isolates were sensitive to Cephalothin while , out of the tested isolates 60% were sensitive to Oxytetracycline 30 mg. Morever, 80% were moderately sensitive to Amoxacillin 30 mg while, 80% of isolates were weakly sensitive to Kanamycin. These results are nearly similar to those reported by Simor and Wilcax (1987) and Mouffok and Lebres (1992).

The antibiogram of C. jejuni isolated from milk and milk products often reflect the misuse of antibiotics in veterinary practice used in virtually all farms may have contributed to the resistance to some antibiotics.

Owing to the importance of C. jejuni as a food-bome pathogen, documented by various outbreaks and sporadic cases of human Campylobacteriosis, allover the world, the importance of carrying out a prospective study on the prevalence of this pathogen in milk and some milk products is widely acknowledged.

References

Abdel-Hady, H. M. (1993): Studies on Campylobacter jejuni and Yersinia enterocolitica as food-poisoning causative organisms in milk and some dairy products. Ph. D. Thesis, Fac. Vet. Med., Cairo Univ., Egypt.

Abdel-Hady, H. M. (1996): DNA prob and test for rapid detection of Campylobacter species in Ovine milk and hard cheese. Vet. Med. J, Giza., 44 (2): 197-201.

Abdel-Samei, H. M. (2000): Campylobacter enteritis associated with milk and its products. A review Article, Dept. of Food Control, Fac. Vet. Med., Moshthor, Zagazig Univ. Egypt.

Allos, B. M. (1997): Association between Campylobacter infection and Guillain-Barre syndrome. J. Infect. Dis., 176: 125-128.

Ansary, A. and Veloo, V.S.L (1991): Conjugal transfer of antibiotic resistance in Campylobacter coli and Campylobacter jejuni isolates of poultry. Tropical Biomedicine, 8 (1) : 77-80.

A. P. H. A. (1992): Standard Methods for the Examination of Dairy Products. 16th Ed. American public Health Association, New York.

Bachmann, H. P. (1994): The fate of potentially pathogenic becteria in Swiss hard and semihard cheeses made from raw milk. J. Dairy Sci., 78: 476-483.

Baron, E. J.: Peterson, L. R. and Finegold, B. M. (1994): Bailey and Scott's Diagnostic Microbiology. 9th Ed., Mosby, st. louis, Baltimore.

Baroun, L. and Ou, J. T. (1991): Strain in expression of virulence by the 90 kilobase pair virulence plasmid of salmonella serovar typhimurium. Microbial. Pathol., 10 : 247 – 251.

Barrett, N. J. (1986): Communicable disease associated with milk and dairy products in England and Wales: 1983 – 1984. J. Infect., 12 : 265-272.

Bean, N. H.; Goulding, J. S.; Lao, C. and Angulo, F. J. (1996): Surveillance of food borne disease outbreaks. United states, 1988-1992. Morbid. Mortal. Weakly Rep. (55-5): I.

Boer, E. DE; Hartog, B. J. and Borst, G. H. A. (1984): Milk as a source of Campylobacter jejuni. Netherlands Milk and Dairy J., 38 (3): 183-194.

Butzler, J. P. and Oosterom, J. (1991): Campylobacter pathogenicity and significance in foods. Int. J. Food Microbiol., 12 (1): 1-8.

Casini, T.; Cristiano, R.; Salvi, G.; Franchini, F.; Calabri, G.; Maria, M.; Mannelli, F.; Salvatore, A.; Doyle, M. P. (1944): A Vibrio associated with swin dysentery. Amer. J. Vet. Res., 5:15.

Doyle, M. P. and Roman, D. J. (1981): Growth and survival of Campylobacter fetus subsp. Jejuni as a function of temperature and pH. J. Food Prot., 44 (8): 596-601.

Doyle, M. P. (1991): Campylobacter jejuni in food-borne diseases. D. O. Cliver. (Ed.), Academic press Inc., PP: 217-222.

DuzGun, S.; Inal, U. and Turk, N. (2000): Isolation of some pathogenic bacteria from the milk samples taken from the milk collection tanks. Sut Taplama Tanklarindanalinan sut orneklerinden Bazi pathojen bakterilerin izolasyonu. Bornova Veteriner Kntorl Ve Arastirma Enstitusu Dergisi., 25 (39): 11-15. Dairy Sci. Abst., 63 (12): 996. 2001.

Edmonds, P.; patton, C. M.; Griffin, P. M.; Barrett, T. J.; Schmid, G. P.; Baker, C. N.; Lambert, M. A. and Brenner, D. J. (1987): Campylobacter hyointestinalis associated with human gastrointestinal disease in the United Stales. J. Clin. Microbiol., 25 : 685-691.

Ehlers, J. G.; Chapparo-serrano, M.; Richter, R. L. and Vanderzant, C. (1982): Survival of Campylobacter fetus subsp. jejuni in Cheddar and Cottage cheese. J. Food Prot., 45 (11): 1018-1021.

EI-Nokrashy, Soheir, A.; EI-Dairouty, R. K.; Effat, B. (1998): Incidence and viability of Campylobacter jejuni in Damiatti cheese. Arab Universities J. Agric. Sci., 6 (2): 471-480. Dairy Sci. Abst., 61 (1) 1999.

EI-Nokrashy, Soheir A.; EI-Magduib, N. and EI-Dairouty, R. K. (1997): Isolation, characterization and thermal inactivation of Campylobacter spp. from Egyptian raw milk. J. MicrobioL, 32 (I): 117-127.

El-Prince, Enas, M.; Hussein, Asmaa, A. and El Said, M. M. (1998): A cohort study of Campylobacter species in dairy cows and infants and their mother's milk in Assiut governorate. 8th Sci. Cong., Fac. Vet. Med., Assiut Univ., Egypt. 33-48.

Ewyda, Elham, H. (1990): Prevalence of Campylobacter spp. in patients with diarrhea in Assiut. M. V. Sc. Thesis, Fac. Med., Assiut Univ., Egypt.

Federighi, M.; Magras, C.; Pilet, M. F.; Wood Word, D.; Johnson, W.; Jugiau, F. and Jauve, J. L. (1999): Incidence of thermotolerant Campylobacter in food. Assessed by Nfiso 10272 Standards: result of two years study. Food MicrobioL, 16:195-204.

Fennell, C. L.; Totten, F. A.; Quinn, J. C.; Patton, D. L.; Molmes, K. K. and Stamm, W. E. (1984): Characterization of Campylobacter-like organisms isolated from homosexual men. J. Infect. Dis., 149:58-66.

Frazier, W. C. and Westhoff, D. C. (1988): Food Microbiology. 4th Ed., Mc Graw Hill Book co., New York.

Hudson, J. A.; Nicol, C.; Wright, J.; Whyte, R. and Hasel, S. K. (1999): Seasonal variation of Campylobacter types from human cases, veterinary cases, raw chicken, milk and water. J. Appl. Microbiol. 87:115-124.

Hudson, S. J.; Sobo, A. 0.; Russel, K. and Ligtfoot, N. F. (1990): Jackdaws as potential source of milk-bome Campylobacter jejuni infection. Lancet (Brit. Ed.), 335:8698,11600.

Humphrey, T. J. and Beckett, P. (1987): Campylobacter jejuni in dairy cows and raw milk. Epidemic!. Infect., 98: (3) 263-269.

Hutchinson, D. N.; Bolton, F. J.; Jelley, W. C. N.; Mathwes, W. G.; Telford, D. R.; Counter, D. E.; Jessop, E. G. and Horsely, S. D. (1985): Campylobacter enteritis associated with consumption of raw goat's milk. Lancet I., 843b:1037-1038.

Inokova, G. and Ivanova, K. (1996): Morphological, cultural and biochemical characteristics of strains of Campylobacter species isolated from faecal and milk samples

in Plovidiv region. Bulgarian J. Agric. ScL, 2 (2): 267-273.

Jay, J. M. (1996): Modem Food Microbiology. 5th Ed., Chapman & Hall, New York.

Kalman, Maria; Szollosi, E.; Czermann, B.; Zimanyi, M.; Sekerer, S. and Kalman, M. (2000): Milk borne Campylobacter infection in Hungary. J. Food Prot., 63 (10): 1426-1429.

Kaufman, P.B.; Wu,W.; Kirn, D. and Csek, L. J.(1995): Handbook of Mollecular and Cellular Methods in Biology and Medicine . CRC Boca, Raton, London.

Lampert, L. M. (1975): Modem Dairy Products. 3rd Ed. , Chemical Publi.Co. Inc., New York, U.S.A.

Lax, A. G.; Barrow, P. A.; Jones, P.W. and Wallis, T. S. (1995): Current perspectives in Salmonellosis, Br. Vet. J., 151:351-377.

Lekowska Kochanial, A; Rozynek, E. and Popowski, J. (1996): Antibiotic resistance of Campylobacter jejuni with reference to plasmid profiles of clinical and chicken isolates. Acta Micro biologica Polonica,45(3-4): 249.

Mentzing, L. O. (1981): Water-borne outbreak of Campylobacter enteritis in central Sweden. Lancet, 5:352-354.

Mouffok, F. and Lebres, E. (1992): Results of technique of isolation and identification of Campylobacter in food. Arch. Inst Pasteur Algerie, (S8):239-246.

Nachamkin, I.; Blaser, M. J. and Tompkins, L. S. (1992): Campylobacter Jejuni. Current strategy and future trends. Amer. Soc. Microbiol. Washington D. C., pp. 3-296.

Nielsen, E, M.; Engberg, J.; Fussing, N.; Petersen. L.; Brogen, C. H, and Stephen, L. W. (2000): Evaluation of phenotypic and genotypic methods for subtyping Campyiobacter jejuni isolates from human, poultry and cattle. J. Clin. Microbiol., PP.: 3800-3810.

Oxoid Manual (1990): Culture media; ingredients and other laboratory services. 6th Ed. Publ. by Oxoid limited, London.

Penner, J.L. (1988): The genus Campylobacter : a decade of progress. Clin. Microbiol.Rev., 1; 157-172.

Phillips, C. A. (1995): Incidence of Campylobacter and possible mode of transmission. NutFood Sci. (I): 12-17.

Ray, B. and Johnson, C. (1984): Survival and growth of freez-stressed Campylobacter jejuni cell in selective media J. Food Safety, 6:183.

Reed, G. H. (1994): Food-bome illness food-bome Campylobacteriosis. Dairy Food and Environ. Sanitation, 14(3): 161-162.

Richter, R. L.; Ledford, R. A. and Murphy, S. C. (1992): Milk and milk products. In: Compendium of Methods for the Microbiological Examination of Foods.3rd Ed. For Venderzant, and D. F. Splittstoesser. (eds-)., American Public Health Association, Washington, D. C. 191-Rosef, 0. and Yundestad, N. (1982): Some characteristic of Campylobacter fetus subsp. jejuni isolated from pigs, birds and man. Acta Vet. Scand., 23: 9-12.

Rosef, O. and Yundestad, N. (1982): Some characteristic of Campylobacter fetus subsp. Jejuni isolated from pigs, birds and man. Acta Vet. Scand., 23 : 9 - 12.

Roshdy, M. SH. (2000): Prevalence of Campylobacterjejuni in raw milk and infantile diarrhea in Assiut. M. V. Sc Thesis, Fac. of med., Assiut Univ., Egypt.

Simor, A. E. and Wilcax, L. (1987): Enteritis associated with Campylobacter laridis. J. Clin. Microbiol., 25 (I): 10-12.

Skirrow, M. B. (1990): Food-home illness. Campylobacter Lancet Octob., 13 (336): 921-924.

Smith, J. L. (1995): Arthritis, Guillain Barre-syndrome, and other sequelae of Campylobacter jejuni enteritis. J. Food Prot., 58:1153-1170.

Smith, S. L; Coker, A. 0. and Olukoyz, D. K. (1997): Biotyping of Campylobacter strains isolated in Lagos Nigeria using the modified preston biotype 2. nature forsch (c), 52, 3-4: 259-263.

Tauxe, R, V. (1992): Epidemiology of Campylobacterjejuni infections in the United States and other industrialized nations. In Campylobacter jejuni: current status and future

trends. American Society of Microbiology, Chap.2, pp.; I. Nachamkin, M. J. Blaser and L. S. Tompkins (eds.), Washington, D. C.

Walker, R. L; Caldwell, M. B.; Lee, E.C.; Gurry, P.; Trust, T. J. and Ruis palacios, G. M. (1986): Pathophysiology of Campylobacter enteritis. Microbiol.

Rev., 50 (I): 81-94.

Waterman, S. C.; Park, R. W. A. and Bramley, A. J. (1984): A search for the source of Campylobacter jejuni in milk. J. Hyg. 93 (2): 333-337.

Wegmuiller, B.; Luthy, J. and Candrian, U.(1993): Direct polymerase chain reaction detection of Campylobacter coli in raw milk and dairy products, Appl. Environ.Microbiol., 59 (7): 2161-2165.

White, C. H. (1986): Occurrence offood-bome pathogens in dairy product. J. Dairy Sci., 69 (Suppl. I): 224.